

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 921 193 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
09.06.1999 Bulletin 1999/23

(51) Int. Cl.⁶: C12N 15/12, C12N 15/24,
C12N 15/88, A61K 48/00

(21) Application number: 98100135.7

(22) Date of filing: 07.01.1998

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 05.12.1997 US 985526

(71) Applicant:
Mixson, Archibald James
Rockville, MD 21201 (US)


(72) Inventor:
Mixson, Archibald James
Rockville, MD 21201 (US)

(74) Representative:
Sleckmann, Ralf, Dr. et al
Cohausz Hannig Dawidowicz & Partner
Patent- und Rechtsanwaltskanzlei
Schumannstrasse 97
40237 Düsseldorf (DE)

(54) Carrier: DNA complexes containing DNA encoding anti-angiogenic peptides and their use in gene therapy

(57) Carrier complexes comprising DNA encoding an anti-angiogenic gene or peptide and optionally a further DNA encoding a tumor suppressor protein are described. When administered to a subject bearing a tumor, the complexes can inhibit growth of the tumor.

Figure 1
Intratumoral Injections of Liposome:DNA Complexes



* Angio vs. BAP, $p < 0.05$

EP 0 921 193 A1

Description

FIELD OF THE INVENTION

- 5 [0001] The present invention relates to delivery of antiangiogenic genes or DNA encoding anti-angiogenic peptides to a tumor in vivo, and expression of the DNA to inhibit tumoral growth. Carrier:DNA complexes are provided comprising DNA encoding at least one anti-angiogenic protein or peptide, optionally together with further DNA encoding a tumor suppressor protein. These complexes are useful in gene therapy for inhibition of tumor growth.

10 BACKGROUND OF THE INVENTIONI. Gene Therapy

- 15 [0002] Development of gene therapy techniques is approaching clinical realization for the treatment of neoplastic and metabolic diseases. There remains substantial need for improvement both in the vector delivery systems for delivery of the transgene to target tissues, and the identification of genes most effective for anti-tumor therapy.

- [0003] Vectors for carrying genes may be viral or non-viral. For example, replication-deficient retroviral vectors can efficiently transfect dividing cells. Local intratumoral injection of retroviruses that contain a thymidine kinase transgene has been used successfully to affect regression of gliomas (Culver et al, *Science*, 256:1550-1552 (1992)). Unlike retro-
20 viral vectors, adenoviral vectors can also transfect non-dividing cells, and their ability to cause insertional mutagenesis is greatly reduced. However, they can have the undesirable potential to activate the immune system in humans (Crystal, *Science*, 270:404-410, (1995). Attempts are underway to minimize the immunogenicity of the adenoviral vectors.

- [0004] Non-viral vectors of DNA include primarily liposomes, peptides, proteins and polymers (Ledley, *Current Opin-
ion in Biotechnology*, 5:626-636 (1994)). Of these, liposomes are currently the most common non-viral vectors of DNA.
25 The major advantage of liposomes over retroviruses is that DNA is not incorporated into the genome, and unlike adenoviral vectors, they are not immunogenic. However, the major limitation of liposomes is that they are not as efficient as viral vectors in transfecting many cell types. Until recently, their medical utility was limited by their rapid uptake by phagocytic cells. Interest in liposomes as a vector has been increased by two technological advances. First, stearily stabilized (Stealth) liposomes have been developed which are more non-reactive and are not readily taken up by the
30 reticuloendothelial system (RES). Stealth liposomes are composed of lipids rich in oxygen in their head group (ethylene glycol or glycolipids) which provide a stearic barrier outside of the membrane. As a result, Stealth liposomes remain in the blood for up to 100 times longer than conventional liposomes, and can thus increase pharmacological efficacy (Papahajopoulos, In: *Stealth Liposomes*, Ed., Lasic et al, CRC Press (1995); and Lasic et al, *Science*, 267:1275-76 (1995)). However, stealth liposomes are still not particularly efficient in transfection of cells or as vectors for DNA.

- 35 [0005] The second significant advance in liposome technology has been the use of cationic liposomes complexed to negatively-charged DNA. Cationic liposomes can condense DNA, and increase transfection yields several orders of magnitude. In the cationic liposome:DNA complex, the nucleic acids or oligonucleotides are not encapsulated, but are simply complexed with small unilamellar vesicles by electrostatic interactions. The exact nature of the cationic liposome:DNA complex is not fully known, but intricate topological rearrangements of the cationic liposome:DNA complex
40 may occur, including DNA condensation, liposome aggregation, and fusion. This supramolecular complex can be added to cells *in vitro*, injected parenterally, or aerosolized for pulmonary applications (Lasic et al, *Science*, 267:1275-1276 (1995)). Further, the intravenous injection into mice of high concentrations of the CAT gene (100 µg or greater) complexed with cationic liposomes has been found to result in 40% transfection efficiency of well vascularized tissues, such as the spleen (Zhu et al, *Science*, 261:209-211 (1993)). Notwithstanding these advances, a major challenge of
45 gene therapy remains the systemic delivery of transgenes to the tumor or peritumoral area that will effectively decrease the size of primary tumors and their metastases. Unlike the spleen and bone marrow, which are highly vascular and have a high capacity to filter macromolecules from the blood stream, most organs and tumors do not have this capacity, and the transfection efficiency of these tissues with liposomes is low (Marshall, *Science*, 269:1051-1055 (1995)). In addition, another limitation of cationic liposome: DNA complexes is that their 1/2 life in the blood stream is normally less
50 than one hour (Allen et al, In: *Liposome Technology*-Vol. III, Ed., Gregoriadis G et al, CRC Press (1993); Li and Huang, *J. of Liposome Research*, 6:589 (1996). Sufficient transfection of the target cell by vectors carrying therapeutic genes has thus far been the rate-limiting step in gene therapy.

II. Tumor Suppressor Genes

- 55 [0006] Tumor suppressor genes are well-known in the art, and include the p53 gene (Baker et al, *Science*, 249:912-915 (1990)), the p21 gene (El-Deiry et al, *Cell*, 75:817-825 (1993); and Harper et al, *Cell*, 75:805-816 (1993)), and the rb gene (Bookstein et al, *Science*, 247:712-715 (1990)).

[0007] Mutations in the tumor suppressor gene p53 are known to occur in over 50% of human tumors, including metastatic breast cancer. Various groups have found that reintroduction of the wild-type P53 by mediated transfer of a single copy of the p53 transgene into a variety of tumor cells, including breast cancer cells, results in a decrease in growth rate and/or attenuated tumor development once those transfected cells were implanted into nude mice (Wang et al, *Oncogene*, 8:279-288 (1993); Baker et al, *Science*, 249:912-915 (1990)); Bookstein et al, *Science*, 247:712-715 (1990); Cheng et al, *Cancer Res.*, 52:222-226 (1992); Isaacs et al, *Cancer Res.*, 51:4716-4720 (1991); Diller et al, *Mol. Cell. Biol.*, 10:5772-5781 (1990); Chen et al, *Oncogene*, 6:1799-1805 (1991); and Zou et al, *Science*, 263:526-529 (1994)). In addition, intratracheal injection of a retrovirus containing the p53 transgene has been shown to significantly inhibit the growth of lung tumors (Fujiwara et al, *J. Natl. Cancer Inst.*, 86:1458-1462 (1994)).

[0008] Systemic intravenous administration of a β -actin promoter-containing vector containing the p53 coding sequence complexed to cationic liposomes has been found to affect the tumor growth of a malignant line of breast cancer cells injected into nude mice (Lesoon-Wood et al, *Proc. Am. Ass. Cancer Res.*, 36:421 (1995); and Lesoon-Wood et al, *Human Gene Ther.*, 6:39-406 (1995)). Of the 15 tumors treated in this study, four of these tumors did not respond to treatment. Because of the unresponsiveness of these tumors, new therapies were sought in the present invention to more effectively decrease the size of these tumors.

[0009] p53 coordinates multiple responses to DNA damage. DNA damage results in an increase in the level of the p53 protein. Following DNA damage, an important function of wild-type p53 function is to control the progression of cells from G1 to S phase. Recently, several groups have found that p53 transcriptionally activates a p21 kd protein (also known as WAF1 or CIP1), an inhibitor of cyclin-dependent kinases (CDKs) (El-Deiry et al, *supra*; and Harper et al, *supra*). Inhibition of CDK activity is thought to block the release of the transcription factor E2F, and related transcription factors from the retinoblastoma protein RB, with consequent failure to activate transcription of genes required for S phase entry (Harper et al, *supra*; and Xiong et al, *Nature*, 366:701-704 (1993)). Evidence consistent with the model that pRb is a downstream effector of p53-induced G1 arrest has recently been reported (Dulic et al, *Cell*, 76:1013-1023 (1994)). Thus, p53 regulates cell cycle through two proteins: p21 and rb.

III. Anti-Angiogenic Proteins

[0010] Proteins with anti-angiogenic activities are well-known and include: thrombospondin I (Kosfeld et al, *J. Biol. Chem.*, 267:16230-16236 (1993); Tolsma et al, *J. Cell Biol.*, 122:497-511 (1993); and Dameron et al, *Science*, 265:1582-1584 (1995)), IL-12 (Voest et al, *J. Natl. Cancer Inst.*, 87:581-586 (1995)), protamine (Ingber et al, *Nature*, 348:555-557 (1990)), angiostatin (O'Reilly et al, *Cell*, 79:315-328 (1994)), laminin (Sakamoto et al, *Cancer Res.*, 5:903-906 (1991)), endostatin (O'Reilly et al, *Cell*, 88: 277-285 (1997)), and a prolactin fragment (Clapp et al, *Endocrinol.*, 133:1292-1299 (1993)). In addition, several anti-angiogenic peptides have been isolated from these proteins (Maione et al, *Science*, 247:77-79 (1990); Woltering et al, *J. Surg. Res.*, 50:245-251 (1991); and Eijan et al, *Mol. Biother.*, 3:38-40 (1991)).

[0011] Thrombospondin I (hereinafter "TSPI") is a large trimeric glycoprotein composed of three identical 180 kd subunits (Lahav et al, *Semin. Thromb. Hemostasis*, 13:352-360 (1987)) linked by disulfide bonds (Lawer et al, *J. Cell Biol.*, 103:1635-1648 (1986); and Lahav et al, *Eur. J. Biochem.*, 145:151-156 (1984)). The majority of anti-angiogenic activity is found in the central stalk region of this protein (Tolsma et al, *supra*). There are at least two different structural domains within this central stalk region that inhibit neovascularization (Tolsma et al, *supra*).

[0012] Besides TSPI, there are six other proteins (fibronectin, laminin, platelet factor-4, angiostatin, endostatin and prolactin fragment) in which peptides have been isolated that inhibit angiogenesis. In addition, the dominant negative fragment of FIK1 and analogues of the peptide somatostatin are known to inhibit angiogenesis.

[0013] Fibronectin (FN) is a major surface component of many normal cells, as well as a potent cell spreading factor. During transformation, the loss of cellular FN has been observed. Furthermore, the addition of fibronectin to transformed cells restores the normal phenotype. It has been found that either heparin-binding or cell-adhesion fragments from FN can inhibit experimental metastasis, suggesting that cell surface proteoglycans are important in mediating the adhesion of metastatic tumor cells (McCarthy et al, *J. Natl. Cancer Inst.*, 80:108-116 (1988)). It has also been found that FN and one of its peptides inhibits *in vivo* angiogenesis (Eijan et al, *Mol. Biother.*, 3:38-40 (1991)).

[0014] Laminin is a major component of the basement membrane, and is known to have several biologically active sites that bind to endothelial and tumor cells. Laminin is a cruciform molecule that is composed of three chains, an A Chain and two B chains. Several sites in laminin have been identified as cell binding domains. These sites promote cellular activities *in vitro*, such as cell spreading, migration, and cell differentiation. Two peptides from two sites of the laminin B1 chain are known to inhibit angiogenesis (Grant et al, *Path. Res. Pract.*, 190:854-863 (1994)).

[0015] Platelet factor-4 (PF4) is a platelet α -granule protein originally characterized by its high affinity for heparin. The protein is released from platelets during aggregation as a high molecular weight complex of a tetramer of the PF4 polypeptide and chondroitin sulfate, which dissociates at high ionic strength. PF4 has several biological properties including immunosuppression, chemotactic activity for neutrophils and monocytes as well as for fibroblasts, inhibition of

bone resorption, and inhibition of angiogenesis. The angiostatic properties of human PF4 are associated with the carboxyl-terminal, heparin binding region of the molecule. A 12 amino acid synthetic peptide derived therefrom has been discovered to have marked angiostatic affects (Maione et al, *Science*, 247:77-79 (1990)).

5 [0016] Endostatin is a 20 kDa protein fragment of collagen XVIII. It has recently been found to be a potent inhibitor of tumor angiogenesis and tumor growth (O'Reilly et al., *Cell*, 88, 277-285, 1997).

[0017] Although somatostatin is not a protein, it is a naturally-occurring cyclic 14 amino acid peptide whose most-recognized function is the inhibition of growth hormone (GH) secretion. Somatostatin is widely distributed in the brain, in which it fulfills a neuromodulatory role, and in several organs of the gastrointestinal tract, where it can act as a paracrine factor or as a true circulating factor. The role played by the neuropeptide somatostatin, also known as somatotropin release inhibitory factor (SRIF), in human cancer is not well understood. Recent investigations involving somatostatin receptors in normal and neoplastic human tissues suggest that the action is complex, and involves both direct and indirect mechanisms. One of the anti-tumor mechanisms of these synthetic somatostatin analogues may be an anti-angiogenic effect (Woltering et al, *J. Surg. Res.*, 50:245-50 (1990)). In a recent study, the ability of native somatostatin and nine somatostatin analogues to inhibit angiogenesis were evaluated. The most potent somatostatin analogue was found to be approximately twice as potent as the naturally-occurring somatostatin (Barrie et al, *J. Surg. Res.*, 55:446-50 (1993)).

[0018] Angiostatin is a 38 kDa polypeptide fragment of plasminogen. Whereas plasminogen has no fibrinogenic activity, angiostatin has marked angiogenic activity (O'Reilly MS, et al *Cell*, 79:315-28 (1994)). Angiostatin was isolated when it was observed that the primary tumor suppressed metastases. That is, when the primary tumor was removed, the metastases grew. Administration of angiostatin blocks neo-vascularization and growth of metastases.

[0019] The Flk1 receptor is a receptor for vascular endothelial growth factor (VEGF). Flk-1 is exclusively expressed on the surface of the endothelial cells. Once VEGF binds to the receptor, the Flk-1 receptor then homodimerizes to stimulate the endothelial cell to divide. If a mutant receptor of Flk-1 is transfected into the endothelial cells, the mutant receptor dimerizes with the wild-type Flk-1 receptor. In this endothelial transfected with the mutant Flk-1 receptor, VEGF is unable to stimulate the endothelial cells to divide. Co-administration of a retrovirus carrying the Flk-1 cDNA (Millauer B. et al., *Nature*, 367, 1994) inhibits tumor growth. This emphasizes that the receptor plays a critical role in the angiogenesis of solid tumors.

[0020] Finally, a 16kd fragment of prolactin has been found to be antiangiogenic. Similar to plasminogen, prolactin is not anti-angiogenic but the prolactin fragment is a potent *in vivo* and *in vitro* inhibitor of angiogenesis (Clapp C. et al. *Endocrinology*. 133:1292-1299 (1993)).

[0021] Despite the evidence that anti-angiogenic peptides can be useful anti-tumor agents, and interest in targeting genes toward the vasculature, there have been no published reports on effective *in vivo* gene therapy regimens utilizing anti-angiogenic DNA sequences.

[0022] The only transfected antiangiogenic gene that has been shown to inhibit tumor growth is full length thrombospondin I. In that study (Weinstat-Saslow et al, *Cancer Research* 54, 6504-6511, (1994)) tumor cells that expressed 15-fold higher levels of the thrombospondin I *in vitro* than baseline cells were implanted into mice. This transfected full length thrombospondin I was secreted from the tumor cells, and effectively reduced the tumor by 60%. Thus, this study determined that transfection of 100% of the tumor cells with a highly expressed and secreted antiangiogenic gene was able to reduce tumor size.

40 SUMMARY OF THE INVENTION

[0023] An object of the invention is to deliver anti-angiogenic genes and/or DNA encoding anti-angiogenic peptides to a tumor site *in vivo*, preferably by injection, whereby the DNA is expressed at the site of the tumor to inhibit tumoral growth.

[0024] A further object of the present invention is to provide carrier complexes containing DNA encoding anti-angiogenic peptides. The carrier may be specifically targeted to the tumor and/or to the tumor vasculature. The complexes are useful for providing anti-angiogenic gene therapy and inhibiting tumor growth in a subject.

50 [0025] A further object of the present invention is to provide carrier complexes containing DNA encoding an anti-angiogenic gene or peptide, or DNA encoding more than one anti-angiogenic gene or peptide, and additionally DNA encoding a tumor suppressor gene.

[0026] In currently preferred embodiments, the carrier material comprises complexes of cationic polymer or cationic liposomes and DNA encoding one or more antiangiogenic peptides, optionally with DNA encoding a tumor suppressor gene.

55 [0027] The complexes are administered in a tumor-inhibiting effective amount to a patient, preferably by injection of the complexes.

[0028] Moreover, a further object of the present invention is the use of DNA encoding at least one anti-angiogenic protein or peptide in a carrier as defined above which is delivered to a tumor site *in vivo* and is expressed at the tumor

site in vivo in the subject for the production of a medicament for inhibiting tumor growth in a subject bearing a tumor, which comprises administering the same to the subject.

[0029] Finally, another object of the present invention is the use of an anti-angiogenic DNA in a form in which the DNA is expressed in the tumor or a peritumoral area inhibiting tumor growth in a subject bearing a tumor, which comprises injecting the same into the subject.

DESCRIPTION OF THE DRAWINGS

[0030]

Figure 1 is a graph depicting the results of the experiment described in Example 8, infra, wherein complexes containing DNA encoding anti-angiogenic peptides were administered intratumorally.

Figure 2 is a graph showing the results of in vitro transfection experiments into endothelial cells using cationic polymer carrier complexed with DNA encoding anti-angiogenic peptides, as described in Example 10.

DETAILED DESCRIPTION OF THE INVENTION

[0031] In one embodiment, the above-described objects of the present invention have been met by a carrier:DNA complex comprising DNA encoding at least one anti-angiogenic gene or peptide and optionally additional DNA encoding a tumor suppressor protein. The DNA may encode a full-length anti-angiogenic protein, or may encode a peptide having antiangiogenic activity, or a combination of DNAs.

[0032] Preferred carrier vehicles are liposomes, polymers, viruses (retroviruses and adenoviruses, for example), viral shells, micelles, microspheres and the like. See, e.g. Nabel, E., Vectors for Gene Therapy, in Current Protocols in Human Genetics on CD-ROM, John Wiley and Sons (1997) The carrier used in the invention is selected such that it can deliver the DNA in vivo to a tumor and/or the peritumoral area, including tumor vasculature, in a manner such that the DNA can be expressed.

[0033] Liposome carriers are known in the art. Reference is made to, for example, Liposome Technology, 2d Edition, CRC Press: Boca Raton (1983); and Stealth Liposomes, Lasic and Martin, Eds., CRC Press: Boca Raton (1995). Examples of cationic lipids include 1,2-dioleoyl-sn-glycero-3-ethylphosphocholine (Avanti, Birmingham, AL), 1,2-dimyristoyl-sn-glycero-3-ethylphosphocholine (Avanti, Birmingham, AL), and (2,3-diol-eyloxy)propyl-N,N,N-trimethyl-ammonium chloride (DOTMA) (Syntex Corp., Palo Alto, CA).

[0034] The cationic lipids may be used in a mixture with dioleoylphosphatidylethanolamine (DOPE) (Avanti, Birmingham, AL). In the cationic liposome embodiment, the amount of cationic lipid present in the mixture is generally in the range of from 100 to 40 mol%, preferably about 50 mol%. The amount of DOPE present in the mixture is generally in the range of from 0 to 60 mol%, preferably about 50 mol%.

[0035] The liposomes may contain lipid derivatives of polyethylene glycol (PEG), referred to herein as "pegylated lipids". Components useful in creating pegylated lipids include, for example, 1,2-diacyl-sn-glycero-3-phosphoethanolamine-N-[poly(ethylene glycol) 2000]. If pegylated lipid components are present, they are generally included in amounts of 0 to 10 mol%, preferably 1 to 5 mol%.

[0036] Cationic liposomes are prepared in a manner similar to other liposomes, for example, the cationic lipids with/without DOPE are dissolved in a solvent, e.g., chloroform. The lipids are then dried in a round bottom flask overnight on a rotary evaporator. The resulting lipids are then hydrated with sterile water over a 1 hr period to form large multilamellar vesicle liposomes. To decrease the size of the liposomes, one may sonicate or pass the liposomes back and forth through a polycarbonate membrane. The DNA is then added to a solution containing the liposomes after their formation.

[0037] Cationic polymer carriers useful in the context of this invention include polyethyleneimine (available from Avanti Lipids), polylysine (available from Sigma), polyhistidine (Sigma), and Superfect (available from Qiagen). Use of cationic polymer carriers for gene delivery in vitro and in vivo has been described in the literature, for example, by Goldman et al., Nature BioTechnology, 15:462 (1997).

[0038] Stealth liposomes concentrate in solid tumors possibly due to their "leaky" vessels. Although stealth liposomes' uptake into cells is decreased due to the pegylation of their surface, this decrease is more than offset by their prolonged half-life in the circulation. Thus, pegylated liposomes are good carriers of DNA. Micelles are closely related to liposomes except they lack a bipolar membrane. They are made up of polar lipids, some of which can be the same cationic lipids utilized in liposomes. Similar to liposomes, micelles are known to transfect cells with plasmid DNA (Zhang YP. Et al., Pharmaceutical Res. 14: 190-6, 1997; Labat-Moleur F. et al., Gene Therapy 3: 1010-7, 1996).

[0039] As known in the art, there are potential problems with the intravenous injection of viral vectors. However, viruses can deliver transgenes by regional intra-arterial and/or intratumoral injections. Construction of viral vector carrying transgenes has been extensively described and they have been used successfully in gene therapy. (Nabel, E. In

„Current Protocols in Human Genetics on CD-ROM“, John Wiley & Sons, Inc. 1997).

[0040] Delivery of the complexes to a target *in vivo* can be enhanced by including a ligand in the complex having affinity for a specific cellular marker. Ligands include antibodies, cell surface markers, viral peptides, and the like, which act to hom the complexes to tumor vasculature or endothelial cells associated with tumor vasculatur , or t tumor cells themselves, if a secreted form of the antiangiogenic DNA is delivered. An antibody ligand may be an antibody or anti-body fragment specific towards a tumor marker such as Her2, CEA, ferritin receptor, or a marker associated with tumor vasculature (integrins, tissue factor, or β -fibronectin isoform). Antibodies or other ligands may be coupled to carriers such as liposomes and viruses, as is known in the art. See, e.g., Neri et al., Nature BioTechnology, 15:1271 (1997); Kir-potin, D. et al., Biochemistry 36:66 (1997) Cheng, Human Gene Therapy, 7:275 (1996); Pasqualini et al., Nature Bio-technology, 15:542 (1997); and Park et al., Proc. Am. Ass. Canc. Res. 38:342 (1997); Mori and Haung *supra*; and Nabel, *supra*. Alternatively, psuedotyping of a retrovirus may be used to target a virus towards a particular cell. Marin et al., Mol. Med. Today, 3:396 (1997).

[0041] In a further embodiment, the complexes further include a tumor suppressor gene. Examples of such tumor suppressor genes include the p53 gene, the p21 gene (El-Deiry et al, *supra*; and Harper, *supra*), and the rb gene (Bookstein et al, *supra*). The p53 gene is the currently preferred tumor suppressor gene.

[0042] The particular anti-angiogenic protein or peptide encoded by the anti-angiogenic DNA is not critical to the present invention. Examples of suitable peptides include:

- (i) a fragment of thrombospondin I (TSPf) having the amino acid sequence shown in SEQUENCE ID NO: 1. This fragment is encoded by the DNA sequence (nucleotides 1013-1650 of the TSPf gene) shown in SEQUENCE ID NO: 2.
- ii) a concatamer of TSPf having the amino acid sequence of SEQUENCE ID NO: 3, which is encoded by the DNA sequence shown in SEQUENCE ID NO: 4.
- iii) laminin peptide having the amino acid sequence shown in SEQUENCE ID NO. 5, which is encoded by the DNA sequence shown in SEQUENCE ID NO. 6.
- iv) a concatamer of the laminin sequence having the amino acid sequence shown in SEQUENCE ID NO: 7, which is encoded by the DNA sequence shown in SEQUENCE ID NO: 8.
- v) a peptide from platelet factor-4 having the amino acid sequence shown in SEQUENCE ID NO: 9, which is encoded by the DNA sequence shown in SEQUENCE ID NO: 10.
- vi) a concatamer of the platelet factor-4 peptide having the amino acid sequence shown in SEQUENCE ID NO: 11, which is encoded by the DNA sequence shown in SEQUENCE ID NO: 12.
- vii) a somatostatin inhibitor peptide having the amino acid sequence shown in SEQUENCE ID NO: 13, which is encoded by the DNA sequence shown in SEQUENCE ID NO: 14.
- viii) a concatamer of somatostatin inhibitor having the amino acid sequence shown in SEQUENCE ID NO: 15, which is encoded by the DNA sequence shown in SEQUENCE ID NO: 16.
- ix) fibronectin inhibitor peptide having the amino acid sequence shown in SEQUENCE ID NO: 17, which is encoded by the DNA sequence shown in SEQUENCE ID NO: 18.
- x) a concatamer of fibronectin inhibitor peptide having the amino acid sequence shown in SEQUENCE ID NO: 19, which is encoded by the DNA sequence shown in SEQUENCE ID NO. 20.
- xi) angiostatin peptide having the amino acid sequence shown in SEQUENCE ID NO: 21, which is encoded by the DNA sequence shown in SEQUENCE ID NO: 22.
- xii) a concatamer of angiostatin peptide having the amino acid sequence shown in SEQUENCE ID NO: 23, which is encoded by the DNA sequence shown in SEQUENCE ID NO: 24.
- xiii) prolactin peptide having the amino acid sequence shown in SEQUENCE ID NO: 25, which is encoded by the DNA sequence shown in SEQUENCE ID NO: 26.
- xiv) a concatamer of prolactin peptide having the amino acid sequence shown in SEQUENCE ID NO. 27, which is encoded by the DNA sequence shown in SEQUENCE ID NO: 28.
- xv) a peptide of Flk-1-DN having the sequence shown in SEQUENCE ID NO: 34, which is encoded by the DNA shown in SEQUENCE ID NO. 35.
- xvi) a peptide of endostatin having the sequence shown in SEQUENCE ID NO: 36, which is encoded by the DNA shown in SEQUENCE ID NO: 37.

[0043] The above sequences are exemplary and not limiting on the scope of the invention. Certain domains of these fragments are known to have antiangiogenic activity, as reported in the literature. As will be apparent, some of these sequences are concatameric. Use of concatamers can increase the anti-angiogenic dosage level without changing the amount of vector necessary for delivery. The concatamers can extend up to approximately 4400 bases in length (the coding region of a large protein), and the number of concatamers possible will depend n the number of bases of a single anti-angiogenic peptide-encoding unit. As seen in the above examples, th concatamer repeats can be separated

by intervening sequences.

[0044] According to a preferred embodiment of the present invention the carrier DNA complexes additionally comprising a marker directing the complexes *in vivo* to a tumor or to tumor or peritumoral area.

5 [0045] Still another preferred embodiment of the present invention is a complex, wherein the DNA is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25; SEQ ID NO:27, SEQ ID NO:31; SEQ ID NO:35; and SEQ ID NO: 37.

[0046] According to a further preferred embodiment of the present invention the tumor suppressor protein used in the aforementioned carrier: DNA complexes is p53.

10 [0047] For fibronectin, the range of concatamers would be about 2 to about 66. Although the maximum number of anti-angiogenic units for the TSPf is about 6, one can increase this concatameric number by deleting sequence less material to anti-angiogenic effects, such as the sequence shown in SEQUENCE ID NO: 29, where the corresponding amino acid sequence is shown in SEQUENCE NO: 30. In a similar manner, the concatameric number of the platelet factor-4 peptide, somatostatin inhibitor, angiostatin, and prolactin can be modified and increased.

15 [0048] Since more than one anti-angiogenic pathway exists, concatamers consisting of two or more types of inhibitor could be more effective than homogenous concatamers. For example, heterogeneous concatamers of TSPI and the fibronectin inhibitors can be inserted into the same vector. An example of such a heterogenous concatamer encoding DNA is shown in SEQUENCE ID NO: 31. In such heterogenous concatamers, the peptide-encoding repeats of each sequence may be linked in blocks and/or randomly.

20 [0049] The heterogeneous concatamers need not be limited to only anti-angiogenic peptides. For example, the protein angiostatin or the large polypeptide fragment of prolactin can be modified with genes encoding anti-angiogenic peptides. Again, the concatameric number will vary depending on the number of nucleotide bases of the unit angiogenic inhibitor. In a concatamer of large and small anti-angiogenic inhibitors, the ratio of large to small inhibitors is 0.1 to 0.9, preferably 1:1.

25 [0050] A translational start signal Met is included in the peptides as well as a transcriptional stop codon (TAA).

[0051] The *SaI* sites present in the above-sequences are a useful cloning tool for insertion of the DNA into a vector, for example BAP vector, which is known to be useful for expressing proteins efficiently *in vivo* from the β -actin promoter (Ray et al, *Genes Dev.*, 5:2265-2273 (1991)). Other restriction sites can be incorporated into the DNA for cloning into other vectors, as those in the art will readily appreciate.

30 [0052] Other useful vectors for containing the DNA sequences include plasmids with a simian viral promoter, e.g., pZeoSV (Invitrogen); the CMV promoter, e.g., pcDNA3, pRc/CMV or pcDNA1 (Invitrogen); or the phosphoglycerate kinase (PGK) promoter (Abud et al., *Developmental Genetics*, 19:51 (1996). Plasmids with a CMV promoter may contain an intron 5' of the multiple cloning site (Zhu et al, *supra*). Plasmids containing the BGH terminator instead of the viral SV40 polyA terminator, e.g., pcDNA3, pRc/CMV, pRc/RSV (Norman et al, IBC's 5th Annual Meeting (1995); and
35 Invitrogen vectors), can also be employed in the present invention so as to increase the expression of the tumor suppressor gene and the anti-angiogenic peptide(s) in targeted cells.

[0053] Expression of the DNA encoding the tumor suppressor protein and the DNA encoding the anti-angiogenic peptide can be achieved using a variety of promoters. For example, the promoter can be a generalized promoter, such as the β -actin promoter, a simian viral promoter, or the CMV promoter, or a tissue specific promoter, such as the α -fetal protein promoter which is specific for liver (Kaneko et al, *Cancer Res.*, 55:5283-5287 (1995), the tyrosinase promoter which is specific for melanoma cells (Hughes et al, *Cancer Res.*, 55:3339-3345 (1995); or the enolase promoter which is specific for neurons (Andersen et al, *Cell. Molec. Neurobiol.*, 13:503-515 (1993)).
40

[0054] The plasmid vector may contain multiple promoters to enhance expression efficiency. Moreover, a plasmid vector may include IRES sequence (internal ribosome entry site) between different DNA coding sequences, allowing for
45 the translation of more than one peptide from the same transcript. Coding sequences can be associated with secretory sequences in the vector to enhance expression levels. In a further embodiment of the invention, the vector may comprise an extrachromosomal replicating vector. See, e.g. Calos, TIG 12:463 (1996). These and other techniques to optimize expression are known to those in the art.

50 [0055] The particular amount of DNA included in the complexes of the present invention is not critical, the amount of total DNA administered in the complexes generally being in the range of about 0.005 to 0.32 μ g/pM of carrier, preferably 0.045 to 0.08 μ g/pM of carrier.

[0056] The DNA encoding a tumor suppressor gene is generally present in an amount of from 0.0025 to 0.16 μ g/pM of carrier, preferably 0.028 to 0.04 μ g/pM of carrier. The DNA encoding an anti-angiogenic peptide is generally present in an amount from 0.0025 to 0.16 μ g/pM of carrier, preferably 0.028 to 0.04 μ g/pM of carrier.

55 [0057] The mole ratio of the DNA encoding the tumor suppressor gene to the DNA encoding the anti-angiogenic peptide is also variable. Generally, the mole ratio is between 1:5 to 5:1, preferably about 1 to 1.

[0058] The DNA encoding the tumor suppressor gene and the anti-angiogenic peptide may be contained on the same vector or on separate vectors. Different DNAs encoding anti-angiogenic peptides may be provided on the same or dif-

ferent vectors within the complexes.

[0059] In another embodiment, the above-described objects of the present invention have been met by a method for inhibiting tumor growth in a subject comprising administering to a tumor-bearing subject a carrier:DNA complex comprising DNA encoding an anti-angiogenic protein or peptide(s) with or without additional DNA encoding a tumor suppressor gene.

[0060] Thus, it refers to the use of DNA encoding at least one anti-angiogenic protein or peptide in a carrier preferably as defined above which is delivered to a tumor site *in vivo* and is expressed at the tumor site *in vivo* in the subject for the production of a medicament for inhibiting tumor growth in a subject bearing a tumor, which comprises administering the same to the subject.

[0061] According to preferred embodiment of the present invention defined in use terms the carrier is selected from the group consisting of liposomes, cationic polymers, micelles, microspheres, viruses, viral components, or combinations of such carriers.

[0062] In a further preferred embodiment the use as instantly claimed further comprises providing a DNA encoding a tumor suppressor protein on the carrier, wherein again preferably the administration is by injection and specifically by intravenous injection.

[0063] Still another object of the present invention is the use of an anti-angiogenic DNA in a form in which the DNA is expressed in the tumor or a peritumoral area inhibiting tumor growth in a subject bearing a tumor, which comprises injecting the same into the subject, wherein preferably DNA encoding a tumor suppressor protein is additionally injected in a form which is expressed in the tumor or associated tumor vasculature.

[0064] The aforementioned injection is preferably done intravenous or intratumoral.

[0065] It is possible to treat different types of tumors. Examples of tumors which can be treated in accordance with the present invention include solid tumors, e.g., lung, colon, brain, breast and melanoma tumors. All of these tumors are very dependent on blood supply to sustain their growth.

[0066] The particular mode of administering the carrier:DNA complex of the present invention depends on various factors, but preferred modes include intravenous, subcutaneous or intratumoral injection. Intravenous injection is the preferred administration mode for distribution of the complex to the developing blood vessels of the tumor.

[0067] The amount of the carrier:DNA to be administered will vary depending upon the age, weight, sex of the subject, as well as the tumor volume and rate of tumor growth in the subject. Generally, the amount of DNA to be administered will be about 1 to 60 μ g, preferably about 5 to 16 μ g.

[0068] The following examples are provided for illustrative purposes and should not be construed as limiting the scope of the invention.

MATERIALS

Production of DNA Vectors

[0069]

A. TSPI Vector

The coding region of the TSPI gene is known (GB Accession code-X14787). The TSPI gene was inserted into the *Xba*I site of BAP vector (Ray et al, *supra*), producing TSPI vector, in which expression of the TSPI gene is controlled by the β -actin promoter.

More specifically, TSPI cDNA and Bluescript plasmid (Promega) were digested with *Hind*I and *Xba*I, and then the TSPI cDNA was ligated into Bluescript. Next, Bluescript containing the TSP cDNA and BAP vector were digested with *Sa*I and *Bam*HI, and TSPI cDNA inserted in the *Xba*I site of BAP vector. The correct orientation of the TSPI gene in BAP vector was confirmed by DNA sequencing.

B. TSPf Vector

TSPf vector is a vector containing a DNA fragment of the TSPI gene which has the two anti-angiogenic domains (nucleotides 992-1650) (Tolsma et al, *supra*), and a start codon and a stop codon. The DNA fragment was prepared by PCR using thrombospondin I cDNA as template, and 100 pmoles of each of the following primers 5'-TAGGTCTAGAATGACTGAAGAGAAACAAAGAG-3' (SEQUENCE ID NO: 32) and 5'-ATGGTCTAGATTAGAGAC-GACTACGTTTCTG-3' (SEQUENCE ID NO: 33) to amplify nucleotides 1013 to 1650 of the TSPI gene. Both primers contain *Xba*I sites (underlined), the first primer contains an ATG start codon (in bold), and the second primer contains a TTA stop codon (reverse orientation in bold).

The resulting 638 base pair fragment of the TSPI gene (hereinafter "TSPf") encodes peptides that are known to be angiogenic inhibitors (Tolsma et al, *supra*).

After amplification, the DNA fragment was purified, digested with *Xba*I, and the digested fragment inserted into

the *Xba*I site of BAP vector such that the expression of the TSPf gene was controlled by the β -actin promoter (Ray et al, *supra*; and Lesoon-Wood et al, *Human Gene Ther.*, 6:395-405 (1995)). The correct orientation of the fragment in BAP vector was verified by digestion with *Bam*HI, and confirmed by DNA sequencing.

5 C. p53 Vector

The coding sequence of the p53 gene was cut from plasmid p1SVhp53c62 (Zakut-Houri et al, *EMBO J.*, 4:1251-1255 (1985)) with *Xba*I, and inserted into the multiple cloning sites of pGEM3Z vector (Promega, Madison, WI). Digestion of the resulting vector with *Sall* and *Bam*HI generated a 1900 bp fragment that was then inserted into the *Sall* and *Bam*HI sites of BAP vector such that expression of the p53 gene was controlled by the β -actin promoter. The correct orientation of the p53 gene in BAP vector was confirmed by DNA sequencing.

D. Laminin peptide vector was prepared by annealing together the following two oligonucleotides:

15 5'-CTATCGTCGACATGTATATTGGTTCTCGTTAAGTCGACCTATC-3' (SEQUENCE ID NO: 38) and
5'-GATAGGTCGACTTAACGAGAACCAATATACATGTCGACGATAG-3' (SEQUENCE ID NO: 39), which contain an anti-angiogenic fragment from laminin, start and stop codons, and *Xba*I restrictions sites. The annealed oligonucleotides were then digested with *Xba*I, and inserted into the *Xba*I site of BAP vector. The plasmid was sequenced to verify correct orientation.

20 E. Angiostatin vector was prepared by amplifying the angiostatin coding sequence of plasminogen cDNA using the following primers:

5'-AGTATCTAGAATGAGTGTATCTGTCACAATG-3' (SEQUENCE ID NO: 40) and
5'-GAATTCTAGATCACCTATGAGGGGTTTGCTC-3' (SEQUENCE ID NO: 41)

25 The resulting amplified fragment, which contained a genetically engineered ATG start site and a TAA stop codon, was digested with *Xba*I, purified, and inserted into the *Xba*I site of BAP vector. The plasmid was sequenced to verify correct orientation.

30 F. Laminin peptide concatamer vector was prepared by initially annealing the following two oligonucleotides:

5' -
CTATCGTCGACATGTATATTGGTTCTCGTAAAAGATATATTGGTTCTCGTGGTAAAAGAGATATTGG
35 TTCTCGTGGTAAAAGATAAGTOGACCTATC-3' (SEQUENCE ID NO: 42)

and 5'-GATAGGTCGACTTAT-3' (SEQUENCE ID NO: 43). The former oligonucleotide contains an anti-angiogenic fragment from laminin repeated four times, start and stop codons, as well as *Xba*I restrictions sites. The annealed oligonucleotides were then extended with PFU (Stratagene), digested with *Sall*, and inserted into the *Sall* site of BAP vector. The plasmid was sequenced to verify correct orientation.

Preparation of Cationic Liposome:DNA Complexes

45 [0070] A DOTMA:DOPE liposome mixture is known to efficiently transfect endothelial cells *in vitro* (Tilkins et al, *Focus*, 16:117-119 (1994)). Accordingly, liposome:DNA complexes were prepared using DOTMA:DOPE, in a 1:1 ratio, essentially as described by Debs et al, *J. Biol. Chem.*, 265:10189-10192 (1990). Similar liposomes preparations can be prepared by mixing, at a 1:1 ratio, DOPE with other cationic lipids, such as, 1,2-dioleoyl-sn-glycero-3-ethylphosphocholine, and 1,2-dimyristoyl-sn-glycero-3-ethylphosphocholine.

50 [0071] More specifically, a mixture of 400 nmoles of the DOTMA and DOPE were dried overnight on a rotary evaporator. Then, the lipids were rehydrated with 1.5 ml of water for 2 hrs. Next, the milky liposome preparation was sonicated with a bath sonicator until clear. The resulting liposome preparation was then passed through a 50 nm polycarbonate filter between 15 to 20 times with a LipsoFast-Basic extruder (Avestin, Ottawa, On).

[0072] The DNA (see following examples) was prepared with the maxi Qiagen kits (Qiagen Inc., Chatsworth, Ca), and washed twice in 70% (v/v) ethanol. The DNA was then washed with distilled water or dialyzed against water for 24 hrs to removed any remaining salt.

[0073] About 400 pmols of the liposome preparation was gently mixed with between 10 to 35 μ g of total DNA in an Eppendorf tube. This amount in each eppendorf tube was sufficient for two injections. The same amount of DNA was

injected in the combination therapies as in the single treatment regimens. For example, if 16 µg of DNA in the combination therapy (8.0 µg of p53 + 8.0 µg of TSPf) was injected into each mouse of one group, then 16 µg of p53 was injected into each mouse of a second group.

5 EXAMPLE 1

[0074] The anti-angiogenic effects of carrier:DNA complexes were evaluated in mice containing MDA-MB-435 breast cancer tumors (American Type Tissue Culture, Bethesda, MD), which are p53 deficient.

10 [0075] More specifically, after administering the anesthetic, Metofane, to 126 female athymic nude mice (NCI), the mice were injected with 2.0×10^5 MDA-MB-435 tumor cells into the mammary fat pad using a stepper (Tridak) and a 27.5 g needle. Two weeks later, the mice whose tumors grew were divided into various treatment regimens, 18 mice per each regimen. The treatment regimens were as follows: (1) untreated; (2) empty BAP vector; (3) TSPI vector alone; (4) TSPf vector alone; (5) p53 vector alone; (6) p53 vector + TSPI vector; and (7) p53 vector + TSPf vector. The mice received two intravenous injections, the first injection 14 days after the malignant cells had been implanted into the mice, and the second injection 24 days after the malignant cells had been implanted into the mice. The first injection consisted of 200 pmoles of the liposomes complexed with 16 µg of total DNA. The second injection consisted of 200 pmoles of the liposomes complexed with 12.0 µg of total DNA. The sizes of the tumors were measured 7 days after the second injection. The results are shown in Table 1 below.

20 TABLE 1

Anti-tumor Effects of TSPI and TSPf	
Putative Anti-tumor DNA	Tumor Size (mm ³)
Untreated	113.5±6.41
BAP	102.9±6.83
TSPI	103.2±8.96
TSPf	89.4±11.06
p53	80.1±12.7*
p53 + TSPI	82.9±6.95*
p53 + TSPf	53.2±8.37**

35 * p53 or p53 + TSPI vs. untreated, $p < 0.05$

** p53 + TSPf vs. untreated or BAP, $p < 0.01$

40 [0076] As shown in Table 1 above, the p53-treated group was found to be statistically different from the untreated group ($p < 0.05$) after 2 injections. However, the p53 treated group was not statistically different from the empty BAP vector group. This was similar to the results described by Lesoon-Wood et al, *Human Gene Ther.*, 6:395-406 (1995), in which p53 was not statistically different from the empty BAP vector group until after 5 injections.

45 [0077] However, p53 in combination with TSPf reduced tumors more effectively than p53 alone. After just 2 injections of this combination therapy, there was a 35% further reduction in tumor growth compared to p53 alone. The combination group was statistically different from both the untreated and the empty BAP vector groups ($p < 0.01$). Although TSPf by itself was slightly less effective than p53, TSPf was, unexpectedly, substantially more effective than TSPI. In fact, the full length TSPI-treatment group had no more effect than either the empty vector or the untreated groups. This was unexpected for several reasons: 1) both the full length and the fragment of thrombospondin I contained the anti-angiogenic peptide, and 2) in a previous ex vivo study (Weinstat-Saslow et al, supra) full length thrombospondin I was effective in inhibiting tumor growth.

EXAMPLE 2

55 [0078] A second experiment was carried out to determine whether the combination therapy of p53 and TSPf was effective at lower dosages, and to confirm that the combination of p53 and TSPf reduced the tumor size significantly more than p53 alone.

[0079] More specifically, 36 mice were injected with 2.0×10^5 MDA-MB-435 tumor cells into the mammary fat pad. Two weeks later, the mice whose tumors grew were divided into various treatment regimens, 12 mice per each regimen.

The treatment regimens were as follows: (1) empty BAP vector; (2) p53 vector alone, and (3) p53 vector + TSPf vector. The mice were injected intravenously with 200 pmols of the liposomes complexed with 8.0 μ g of total DNA. Subsequently, the mice were treated in the same manner with 200 pmols of the liposomes complexed with 12 μ g of total DNA for the next 4 injections. Ten days elapsed between each injection. The sizes of the tumors were measured before each injection and 7 days after the last injection. The results are shown in Table 2 below:

TABLE 2

Anti-tumor Effects of p53 and TSPf	
Putative Anti-tumor DNA	Tumor Size (mm ³)
BAP	855 \pm 345
p53	616 \pm 142
p53 + TSPf	265 \pm 133*

* p53 + TSPf vs. BAP, $p < 0.02$

[0080] As shown in Table 2 above, the combination therapy with p53 and TSPf was statistically different from BAP, whereas the p53 alone treatment was not. This experiment confirmed that p53 and TSPf can be more effective than p53 alone. Furthermore, a different dosage regimen, without an initial booster dose of 16 μ g of DNA as used in the experiment in Table 1, accentuated the difference between the combination treatment and the p53 alone treatments.

EXAMPLE 3

[0081] The experiment of Example 2 was repeated to confirm that BAP-TSPf complexed to liposomes effectively inhibited the growth of implanted tumors. Five injections of the liposome:DNA complex was administered intravenously to three groups: 1) BAP, 2) TSPf, or 3) p53. Results are shown in Table 3.

TABLE 3

Antitumor Effects of TSPf	
Putative Anti-tumor genes	Tumor Size (mm ³)
BAP	619 \pm 65
TSPf	386 \pm 35*
P53	419 \pm 26*

*TSPf vs. BAP $p < 0.05$

p53 vs. BAP, $p < 0.05$

After five intravenous injections at a dose of 14.5 μ g, the TSPf treatment group was statistically different from the BAP group.

EXAMPLE 4

[0082] An experiment was carried out to investigate the efficacy of complexes carrying DNA encoding anti-angiogenic peptide fragments of angiostatin and laminin.

[0083] 126 mice injected with MDA-MB-435 tumor cells as described in Example 2 were treated as follows: (1) BAP vector; (2) TSPf vector alone; (3) laminin peptide vector alone; and (4) angiostatin vector alone. The mice received 4 intravenous injections, the first injection being 10 days after the malignant cells had been implanted into the mice, and the remaining injections were thereafter 10 days apart. The injections consisted of 200 pmols of the liposomes complexed with 12.5 μ g of total DNA.

[0084] The results are shown in Table 4 below.

TABLE 4

Putative anti-tumor DNA	Tumor Size (mm ³)
BAP	194.7 ± 11.9
TSPf	135.9 ± 11.9*
Laminin peptide	126.4 ± 8.4*
Angiostatin	95.2 ± 6.3*,**

* TSP, Laminin peptide, and Angiostatin vs. BAP, p<0.05

** Angiostatin vs. BAP, p<0.01

[0085] As shown in Table 4 above, the cationic liposomes containing DNA encoding anti-angiogenic peptides (TSPf, laminin peptide and angiostatin) significantly inhibited tumor growth.

EXAMPLE 5

[0086] MCF7 cells (American Type Tissue Culture, Bethesda, MD), which are a breast cancer cell line with two normal p53 alleles, were evaluated as described above except that 4.0×10^6 cells were injected into the mice and the third injection contained 12 µg of the DNA. Each injection was 10 days apart. Nine mice were injected with each of the following treatments except for regimen (1), in which 8 mice were treated: (1) untreated; (2) BAP; (3) p53; and (4) p53 + TSP. The sizes of the tumors were measured 7 days after the third injection. The results are shown in Table 5 below.

TABLE 5

Effect of p53 and TSP. on MCF7s Cells	
Putative Anti-tumor Genes	Tumor Size (mm ³)
Untreated	124.6±7.3
BAP	136±16.8
p53	83.1±13.6*
p53 + TSPf	69.0±13.**

* p53 vs. untreated or BAP, p<0.05

** p53 + TSPF vs. untreated or BAP, p<0.01

[0087] As shown in Table 5 above, the most effective therapy against MCF7 was p53 and TSPf. The significance level for the p53 + TSPf therapy was greater than for p53 alone when they were compared against either the untreated or the BAP groups. The above experiment confirmed that p53 and TSPf can decrease the MCF7 tumor more than the p53 treated or the untreated groups.

EXAMPLE 6

[0088] 4×10^5 MCF7 cells were injected bilaterally into the mammary fat pads of the 28 nude mice. After two weeks of growth, these mice were randomly divided into four groups: 1) empty vector, 2) p53, 3) p53 + TSPf, and 4) untreated. The mice received one injection of 200 pmoles of liposomes complexed with 14 µg of DNA, and the tumors from the various treatment groups were measured 10 days after the treatment. The results are shown in Table 6 below.

TABLE 6

Putative Anti-tumor Genes	Tumor Siz (mm ³)
Empty vector-	54.7±4.0
p53	45.5±5.0
p53 + TSPf	33.9±3.6*
Untreated	61.9±8.3

*, p53 + TSPf vs Untreated, $p < .025$

[0089] As shown in Table 6 and previous tables, the additional reduction of the tumor by the combined use of p53 and TSPf compared to the use of p53 only, suggests that TSPf and p53 have different mechanisms of action. Although this does not preclude that the target of p53 is the vasculature of the tumor, the mechanism of inhibition of the tumor by p53 is uncertain at present. However, any mechanism of tumor inhibition by p53 and/or thrombospondin I must account for the low transfection efficiency of the tumor. Using a liposome complexed to a chloramphenicol acetyltransferase marker, it has been demonstrated that less than 5% of the tumor derived from MDA-MB-435 cells was transfected with the marker gene, and assuming similar transfection efficiency here, these favorable results were observed notwithstanding the very low level of transfection.

EXAMPLE 7

[0090] In a further experiment, it was determined that liposomes complexed to DNA encoding the laminin peptide can inhibit tumor growth. More specifically, after administering the anesthetic, Metofane, to 24 female athymic nude mice, the mice were injected with 3.0×10^5 MDA-MB-435 tumor cells into the mammary fat pad using a stepper and a 27.5 g needle. Two weeks later, the mice whose tumors grew were divided into various treatment regimens, 8 mice per each regimen. The treatment regimens were as follows: (1) BAP, (2) laminin, and (3) p53 + laminin. The mice were injected intravenously with 200 pmols of the liposomes complexed with 12.5 μ gs of total DNA 6.25 μ g of each vector when a combination was used. the mice then received 3 injections, each 10 days apart. The tumors were measured at the time of each injection and at the time of the last injection. The results are shown in Table 7 below.

TABLE 7

Putative Anti-tumor Genes	Tumor Size (mm ³)
BAP	345 ± 23.5
Laminin peptide	280 ± 32.4
Laminin peptide + p53	192 ± 10.5*

*BAP vs. Laminin peptide + p53, $p < 0.05$

[0091] As shown in Table 7 above, cationic liposomes containing a combination of DNAs encoding laminin peptide + p53 were more effective in reducing tumor growth than when DNA encoding the anti-angiogenic peptide was used alone. Thus, the addition of a tumor suppressor gene, p53, can enhance the anti-tumor effect of the anti-angiogenic peptide.

EXAMPLE 8

[0092] Although intravenous injection is preferred, the method of administration of the liposome:DNA complex is not critical. It has been found that intratumoral injections are effective. In an experiment involving intratumoral injection, 18 mice were injected with 3×10^5 C6 glioma cells (rat brain tumors) subcutaneously. Six days after the injections, the mice were separated into 3 groups: 1) BAP, 2) FLK-DN (a dominant negative receptor), and 3) angiostatin. After the second intratumoral injection, there was a statistical difference between the angiostatin and the BAP groups. See Figure 1.

Thus, the therapy of the invention is effective when complexes are administered intratumorally. The therapy is effective against tumors other than breast tumors.

EXAMPLE 9

5

[0093] It was also found that a liposome: secretory angiostatin construct can be more effective than the non-secreted analog. In this experiment, 24 nude mice were injected with 3×10^5 MDA-MB-435 cells. Two weeks later the mice were divided into three groups, and received the following therapies intravenously: 1) liposome:BAP, 2) liposome:secreted angiostatin, and 3) liposome:angiostatin. The concentration of DNA injected into the mice was 14.5 ugs. The mice
10 received one injection of the liposome:DNA complex and their tumors were measured 10 days after the injection.

Table 8

15

Efficacy of Secretory Angiostatin	
Therapeutic DNA	Tumor Size (mm ³)
Angiostatin	28.8±2.2
Angiostatin-Secretory	18.6±1.8*
BAP	30.5±3.3

20

*, P<0.05, BAP vs. Angiostatin-secretory

[0094] As seen in table 8, the secretory angiostatin treatment group was more effective than the vector control or the
25 angiostatin treatment group in reducing the size of the tumor. From this experiment, it is demonstrated that a secretory sequence inserted into the 5' portion of the antiangiogenic inhibitor can increase its efficacy.

EXAMPLE 10

30 [0095] Further experiments indicate that cationic polymers can be useful as carriers in the present therapy, and can be the carriers of choice under certain conditions.

[0096] In the following example, a cationic polymer (Superfect) was compared to cationic liposomes as carrier for transfecting endothelial cells *in vitro* with the CAT marker. The cationic liposomes used for comparison to the polymer were DOSPER (Boehringer), which of 14 lipids screened *in vitro* gave the best results. In this experiment, 1×10^6 Huvec
35 cells were placed into each well of a 6 well plate. 25 uis of Superfect complexed with 2 ugs of DNA was added to each plate 24 hours after the initial seeding of the cells, and compared to plates to which had been added 2 ugs of DNA complexed with cationic liposomes. 36 hours after the transfection, the cells were lysed and the amount of CAT protein was assayed. The results are shown in Table 9.

40

Table 9

Vectors	Activity (DPMs/protein)
Cationic liposomes with BAP	31.1±7.2
Cationic liposomes with CAT	682±129
Superfect with BAP	21.4±0.458
Superfect with CAT	10816±687*
p<0.001, Superfect-CAT vs. Cationic liposome-CAT	

50

[0097] This experiment suggests that a cationic polymer are superior in the transfection of endothelial cells, which is significant since we have hypothesized that endothelial cells of the tumor are a primary target of the therapeutic gene. Similarly, it has been found in some cell lines that Superfect is a better transfection agent *in vitro* than cationic lipo-
55 somes.

EXAMPLE 11

[0098] Since Superfect appeared to be superior to cationic liposomes in the transfection of endothelial cells *in vitro*, Superfect complexed to a therapeutic gene was tested to examine if it would inhibit tumor growth more than the corresponding liposome complex. This experiment was based on the hypothesis that the endothelial cells and not other cells are the primary target of these cationic vehicle:DNA complexes. In this experiment, six mice were injected with 2.5×10^5 MDA cells into the mammary fat pad bilaterally. These tumors were allowed to grow to a large size for 2 months. At this size, the tumor growth is rapidly increasing at an exponential rate and is more resistant to treatment compared to smaller tumors. The mice were treated intravenously via the tail vein with either the cationic liposome:BAP-p53/CMV-TSPf or Superfect:BAP-p53/CMV-TSPf. 9.5 μ g of DNA were complexed to Superfect (108 μ g) or the cationic liposome (200 pmoles). The mice received only one dose of these therapies and their tumors were measured 10 days later. The mice tolerated both therapies without any apparent toxicity. The results are given in the table below.

TABLE 10

	Liposome	Superfect
Before Treatment	380 \pm 95 [#]	384 \pm 86
After Treatment	525 \pm 80	403 \pm 72

[#] - tumor size in mm³

The Superfect carrier appears to be superior to the liposome carrier even after one dose in these large tumors. There was only a minimal increase (5%) in the Superfect-treated group whereas there was a marked increase in the liposome-treated group (38%). When the growth of individual tumors was examined and compared to pre-treatment measurements, all 6 tumors in the liposome-treated group increased in their size. In contrast, 4 of the 6 tumors in the Superfect group showed regression in their size compared to pre-treatment measurements.

EXAMPLE 12

[0099] This experiment was carried out using concatamer DNA encoding anti-angiogenic peptides. Mice injected with MDA-MB-435 tumor cells were treated as follows:

(1) BAP vector; (2) laminin peptide concatamer alone; and (3) laminin peptide vector alone. The mice received 2 intravenous injections, the first injection being 10 days after the malignant cells had been implanted into the mice, and the second injection 10 days later. The injections consisted of 200 pmols of liposomes complexed with 12.5 μ g of vector DNA. The results are shown in Table 10 below.

TABLE 11

Putative Anti-tumor DNA	Tumor Size (mm ³)
BAP	86.8 \pm 12.0
Laminin peptide concatamer	63.9 \pm 4.8
Laminin peptide	53.7 \pm 3.0*

* Laminin peptide v. BAP, $p < 0.05$

[0100] As shown in Table 10, complexes containing DNA encoding laminin concatamer or laminin peptide reduced tumor growth compared to the control (BAP vector).

EXAMPLE 13

[0101] To assess efficacy using a combination of DNAs encoding antiangiogenic peptides, mice injected with MDA-MB-435 tumor cells were treated as follows:

[0102] (1) BAP vector; (2) TSPf vector + angiostatin vector; (3) laminin peptide vector + TSPf vector; (4) laminin peptide vector + angiostatin vector; and (5) laminin peptide and FIK-DN receptor. The mice received 5 intravenous injections.

tions, the first injection being 10 days after the malignant cells had been implanted into the mice, and the remaining injections 10 days apart. The injections consisted of 200 pmols of liposomes complexed with 12.5 μg of total vector DNA, with 6.25 μg of each vector when a combination was used. The results are shown in Table 11 below.

5

TABLE 12

10

Putative anti-tumor DNA	Tumor Size (mm^3)
BAP	626 ± 78
TSPf + Angiostatin	$296 \pm 40^*$
Laminin peptide + TSPf	461 ± 54
Laminin peptide + Angiostatin	483 ± 46
Laminin Peptide and FIK-DN	482 ± 21

15

* TSPF + Angiostatin vs. BAP, $p < 0.01$

[0103] As shown in Table 11, cationic liposomes containing combinations of DNA encoding anti-angiogenic peptides showed favorable inhibition of tumor growth.

20 [0104] While the invention has been described in detail and by reference to specific embodiments thereof, it will be apparent to one of ordinary skill in the art that various changes and modifications can be made therein without departing from the spirit and scope thereof. Documents cited herein are incorporated by reference to the extent relevant to practicing the invention.

25

30

35

40

45

50

55

EP 0 921 193 A1

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: Archibald James MIXSON
(B) STREET: 1 Baederwood Ct.
(C) CITY: Rockville
(D) STATE: Maryland
(E) COUNTRY: USA
(F) POSTAL CODE (ZIP): 21201

(ii) TITLE OF INVENTION: CARRIER:DNA COMPLEXES CONTAINING DNA
ENCODING ANTI-ANGIOGENIC PEPTIDES AND THEIR USE IN GENE
THERAPY

(iii) NUMBER OF SEQUENCES: 43

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(v) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: EP 98
(B) FILING DATE: 07.01.1998

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 218 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Met	Thr	Glu	Glu	Asn	Lys	Glu	Leu	Ala	Asn	Glu	Leu	Arg	Arg	Pro	Pro	1	5	10	15
Leu	Cys	Tyr	His	Asn	Gly	Val	Gln	Tyr	Arg	Asn	Asn	Glu	Glu	Trp	Thr	20	25	30	
Val	Asp	Ser	Cys	Thr	Glu	Cys	His	Cys	Gln	Asn	Ser	Val	Thr	Ile	Cys	35	40	45	
Lys	Lys	Val	Ser	Cys	Pro	Ile	Met	Pro	Cys	Ser	Asn	Ala	Thr	Val	Pro	50	55	60	
Asp	Gly	Glu	Cys	Cys	Pro	Arg	Cys	Trp	Pro	Ser	Asp	Ser	Ala	Asp	Asp	65	70	75	80
Gly	Trp	Ser	Pro	Trp	Ser	Glu	Trp	Thr	Ser	Cys	Ser	Thr	Ser	Cys	Gly				

EP 0 921 193 A1

85 90 95

Asn Gly Ile Gln Gln Arg Gly Arg Ser Cys Asp Ser Leu Asn Asn Arg
100 105 110

Cys Glu Gly Ser Ser Val Gln Thr Arg Thr Cys His Ile Gln Glu Cys
115 120 125

Asp Lys Arg Phe Lys Gln Asp Gly Gly Trp Ser His Trp Ser Pro Trp
130 135 140

Ser Ser Cys Ser Val Thr Cys Gly Asp Gly Val Ile Thr Arg Ile Thr
145 150 155 160

Asn Leu Cys Ser Pro Ser Pro Gln Met Asn Gly Lys Pro Cys Glu Gly
165 170 175

Arg Glu Ala Glu Thr Lys Ala Cys Lys Lys Asp Ala Cys Pro Ile Asn
180 185 190

Gly Gly Trp Gly Pro Trp Ser Pro Trp Asp Ile Cys Ser Val Thr Cys
195 200 205

Gly Gly Gly Val Gln Lys Arg Ser Arg Leu
210 215

(2) INFORMATION FOR SEQ ID NO:2:

(1) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 657 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATGACTGAAG AGAACAAAGA GTTGCCAAT GAGCTGAGGC GGCCTCCCCT ATGCTATCAC 60

AACGGAGTTC AGTACAGAAA TAACGAGGAA TGGACTGTTG ATAGCTGCAC TGAGTGTCAC 120

TGTCAGAACT CAGTTACCAT CTGCAAAAAG GTGTCTGCC CCATCATGCC CTGCTCCAAT 180

GCCACAGTTC CTGATGGAGA ATGCTGTCTT CGCTGTTGGC CCAGCGACTC TCGGACGAT 240

GGCTGGTCTC CATGGTCCGA GTGGACCTCC TGTTCTACGA GCTGTGGCAA TGGAAATCAG 300

CAGCGCGGCC GCTCCTGCCA TAGCCTCAAC AACCGATGTG AGGGCTCCTC GGTCCAGACA 360

CGGACCTGCC ACAFTCAGGA GTGTGACAAA AGATTTAAAC AGGATGGTGG CTGGAGCCAC 420

TGGTCCCCGT GGTCTCTTTG TTCTGTGACA TGTGGTGATG GTGTGATCAC AAGGATCCGG 480

CTCTGCAACT CTCCCAGCCC CCAGATGAAT GGGAAACCCT GTGAAGGCGA AGCGCGGGAG 540

ACCAAAGCCT GCAAGAAAGA CGCCTGCCCC ATCAATGGAG GCTGGGGTCC TTGGTCACCA 600

TGGGACATCT GTTCTGTCAC CTGTGGAGGA GGGGTACAGA AACGTAGTCG TCTCTAA 656

(2) INFORMATION FOR SEQ ID NO:3:

55

EP 0 921 193 A1

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 441 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

5

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

15

20

25

30

35

40

45

50

55

Met Thr Glu Glu Asn Lys Glu Leu Ala Asn Glu Leu Arg Arg Pro Pro
 1 5 10 15
 Leu Cys Tyr His Asn Gly Val Gln Tyr Arg Asn Asn Glu Glu Trp Thr
 20 25 30
 Asp Val Ser Cys Thr Glu Cys His Cys Gln Asn Ser Val Thr Ile Cys
 35 40 45
 Lys Lys Val Ser Cys Pro Ile Met Pro Cys Ser Asn Ala Thr Val Pro
 50 55 60
 Asp Gly Glu Cys Cys Pro Arg Cys Trp Pro Ser Asp Ser Ala Asp Asp
 65 70 75 80
 Trp Gly Ser Pro Trp Ser Glu Trp Thr Ser Cys Ser Thr Ser Cys Gly
 85 90 95
 Gly Asn Ile Gln Gln Arg Gly Arg Ser Cys Asp Ser Leu Asn Asn Arg
 100 105 110
 Cys Glu Gly Ser Ser Val Gln Thr Arg Thr Cys His Ile Gln Glu Cys
 115 120 125
 Asp Lys Arg Phe Lys Gln Asp Gly Gly Trp Ser His Trp Ser Pro Trp
 130 135 140
 Ser Ser Cys Ser Val Thr Cys Gly Asp Gly Val Ile Thr Arg Ile Thr
 145 150 155 160
 Leu Cys Asn Ser Pro Ser Pro Gln Met Asn Gly Lys Pro Cys Glu Gly
 165 170 175
 Glu Ala Arg Glu Thr Lys Ala Cys Lys Lys Asp Ala Cys Pro Ile Asn
 180 185 190
 Gly Gly Trp Gly Pro Trp Ser Pro Trp Asp Ile Cys Ser Val Thr Cys
 195 200 205
 Gly Gly Gly Val Gln Lys Arg Ser Arg Leu Cys Val Asp Ser Arg Met
 210 215 220
 Thr Glu Glu Asn Lys Glu Leu Ala Asn Glu Leu Arg Arg Pro Pro Leu
 225 230 235 240
 Cys Tyr His Asn Gly Val Gln Tyr Arg Asn Asn Glu Glu Trp Thr Val
 245 250 255
 Asp Ser Cys Thr Glu Cys His Cys Gln Asn Ser Val Thr Ile Cys Lys
 260 265 270

EP 0 921 193 A1

Lys Val Ser Cys Pro Ile Met Pro Cys Ser Asn Ala Thr Val Pro Asp
 275 280 285
 Gly Glu Cys Cys Pro Arg Cys Trp Pro Ser Asp Ser Ala Asp Asp Gly
 290 295 300
 Trp Ser Pro Trp Ser Glu Trp Thr Ser Cys Ser Thr Ser Cys Gly Asn
 305 310 315 320
 Gly Ile Gln Gln Arg Gly Arg Ser Cys Asp Ser Leu Asn Asn Arg Cys
 325 330 335
 Glu Gly Ser Ser Val Gln Thr Arg Thr Cys His Ile Gln Glu Cys Asp
 340 345 350
 Lys Arg Phe Lys Gln Asp Gly Gly Trp Ser His Trp Ser Pro Trp Ser
 355 360 365
 Ser Cys Ser Val Thr Cys Gly Asp Gly Val Ile Thr Arg Ile Thr Leu
 370 375 380
 Cys Asn Ser Pro Ser Pro Gln Met Asn Gly Lys Pro Cys Glu Gly Glu
 385 390 395 400
 Ala Arg Glu Thr Lys Ala Cys Lys Lys Asp Ala Cys Pro Ile Asn Gly
 405 410 415
 Gly Trp Gly Pro Trp Ser Pro Trp Asp Ile Cys Ser Val Thr Cys Gly
 420 425 430
 Gly Gly Val Gln Lys Arg Ser Arg Leu
 435 440

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1326 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ATGACTGAAG AGAACAAAGA GTTGGCCAAT GAGCTGAGGC GGCCTCCCCT ATGCTATCAC 60
 AACGGAGTTC AGTACAGAAA TAACGAGGAA TGGACTGTTG ATAGCTGCAC TGAGTGTCAC 120
 TGTCAAGACT CAGTTACCAT CTGCAAAAAG GTGTCTGCC CCATCATGCC CTGCTCCAAT 180
 GCCACAGTTC CTGATGGAGA ATGCTGTCCT CGCTGTTGGC CCAGCGACTC TGCGGACGAT 240
 GGCTGGTCTC CATGGTCCGA GTGGACCTCC TGTCTACGA GCTGTGGCAA TGGAAATCAG 300
 CAGCGCGGCC GCTCCTGCGA TAGCCTCAAC AACCGATGTG AGGGCTCCTC GGTCCAGACA 360
 CGGACCTGCC ACATTCAGGA GTGTGACAAA AGATTTAAAC AGGATGGTGG CTGGAGCCAC 420
 TGGTCCCCGT GGTCACTCTG TTCTGTGACA TGTGGTGATG GTGTGATCAC AAGGATCCGG 480

EP 0 921 193 A1

5 CTCTGCAACT CTCCCAGCCC CCAGATGAAT GGGAAACCT GTGPAAGCGA AGTCCGGGAG 540
 ACCAAAGCCT GCAAGAAAGA CGCCTGCCCC ATCAATGGAG GCTGGGGTCC TTGGTCACCA 600
 TGGGACATCT GTTCTGTCAC CTGTGGAGGA GGGGTACAGA AACGTAGTCG TCTCTGCGTC 660
 GACTCTAGAA TGAAGAAGA GAACAAAGAG TTGGCCAATG AGCTGAGGCG GCCTCCCCTA 720
 TGCTATCACA ACGGAGTTCA GTACAGAAAT AACGAGGAAT GGACTGTTGA TAGCTGCACT 780
 10 GAGTGTCACT GTCAGAACTC AGTTACCATC TGCAAAAAGG TGTCTGCCC CATCATGCCC 840
 TGCTCCAATG CCACAGTTCC TGATGGAGAA TGCTGTCTC GCTGTTGGCC CAGCGACTCT 900
 GCGGACGATG GCTGGTCTCC ATGGTCCGAG TGGACCTCCT GTTCTACGAG CTGTGGCAAT 960
 15 GGAATTCAGC AGCGCGGCCG CTCCTGCGAT AGCCTCAACA ACCGATGTGA GGGCTCCTCG 1020
 GTCCAGACAC GGACCTGCCA CATTGAGGAG TGTGACAAA GATTAAACA GGATGGTGGC 1080
 TGGAGCCACT GGTCCCCGTG GTCATCTTGT TCTGTGACAT GTGGTGATGG TGTGATCACA 1140
 20 AGGATCCGGC TCTGCAACTC TCCCAGCCCC CAGATGAATG GGAAACCCTG TGAAGGCGAA 1200
 GCGCGGGAGA CCAAAGCCTG CAAGAAAGAC GCCTGCCCCA TCAATGGAGG CTGGGGTCCT 1260
 TGGTCACCAT GGGACATCTG TTCTGTCACC TGTGGAGGAG GGTACAGAA ACGTAGTCGT 1320
 25 CTCTAA 1326

(2) INFORMATION FOR SEQ ID NO:5:

30 (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Tyr Ile Gly Ser Arg
 1 5

40 (2) INFORMATION FOR SEQ ID NO:6:

45 (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GTCGACATGT ATATTGGTTC TCGTTAAGTC GAC 33

(2) INFORMATION FOR SEQ ID NO:7:

55

5 (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

10 (x1) SEQUENCE DESCRIPTION: SEQ ID NO:7:
 Met Tyr Ile Gly Ser Arg Gly Lys Ser Tyr Ile Gly Ser Arg Gly Lys
 1 5 10 15
 Ser Tyr Ile Gly Ser Arg Gly Lys Ser
 20 25

15 (2) INFORMATION FOR SEQ ID NO:8:
 (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 90 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

20 (x1) SEQUENCE DESCRIPTION: SEQ ID NO:8:
 GTCGACATGT ATATTGGTTC TCGTGTA AAA GTTATATTGG TTCTCGTGGT AAAAGTTATA 60
 TTGGTTCTCG TGGTAAAAGT TAAGTCGACC 90

25 (2) INFORMATION FOR SEQ ID NO:9:
 (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 13 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

30 (x1) SEQUENCE DESCRIPTION: SEQ ID NO:9:
 Met Leu Tyr Lys Lys Ile Ile Lys Lys Leu Leu Glu Ser
 1 5 10

35 (2) INFORMATION FOR SEQ ID NO:10:
 (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 54 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

40 (x1) SEQUENCE DESCRIPTION: SEQ ID NO:10:
 Met Leu Tyr Lys Lys Ile Ile Lys Lys Leu Leu Glu Ser
 1 5 10

45 (2) INFORMATION FOR SEQ ID NO:11:
 (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 54 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

50 (x1) SEQUENCE DESCRIPTION: SEQ ID NO:11:
 Met Leu Tyr Lys Lys Ile Ile Lys Lys Leu Leu Glu Ser
 1 5 10

55 (2) INFORMATION FOR SEQ ID NO:12:
 (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 54 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:12:
 Met Leu Tyr Lys Lys Ile Ile Lys Lys Leu Leu Glu Ser
 1 5 10

GTCGACATGC TTTATAAGAA GATCATCAAG AAGCTTCTTG AGAGTTAAGT CGAC

54

(2) INFORMATION FOR SEQ ID NO:11:

5

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 46 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

15

Met Leu Tyr Lys Lys Ile Ile Lys Lys Leu Leu Glu Ser Gly Lys Ser
1 5 10 15

Leu Tyr Lys Lys Ile Ile Lys Lys Leu Leu Glu Ser Gly Lys Ser Leu
20 25 30

20

Tyr Lys Lys Ile Ile Lys Lys Leu Leu Glu Ser Gly Lys Ser
35 40 45

(2) INFORMATION FOR SEQ ID NO:12:

25

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 153 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

35

GTCGACATGC TTTATAAGAA GATCATCAAG AAGCTTCTTG AGAGTGGTAA AAGTCTTTAT 60

AAGAAGATCA TCAAGAAGCT TCTTGAGAGT GGTAAAAGTC TTTATAAGAA GATCATCAAG 120

AAGCTTCTTG AGAGTGGTAA AAGTTAAGTC GAC 153

(2) INFORMATION FOR SEQ ID NO:13:

40

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

50

Met Phe Cys Tyr Trp Lys Val Cys Trp
1 5

(2) INFORMATION FOR SEQ ID NO:14:

55

5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:
 GTCGACATGT TCTGTTATTG GAAGGTTTGT TGGTAAGTCG AC 42

15 (2) INFORMATION FOR SEQ ID NO:15:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 34 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:
 25 Met Phe Cys Tyr Trp Lys Val Cys Trp Gly Lys Ser Phe Cys Tyr Trp
 1 5 10 15
 Lys Val Cys Trp Gly Lys Ser Phe Cys Tyr Trp Lys Val Cys Trp Gly
 20 25 30
 30 Lys Ser

(2) INFORMATION FOR SEQ ID NO:16:
 35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 117 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:
 45 GTCGACATGT TCTGTTATTG GAAGGTTTGT TGGGGTAAAA GTTCTCTGTTA TTGGAAGGTT 60
 TGTGTTGGGTA AAAGTTTCTG TTATTGGAAG GTTTGTTGGG GTAAAAGTTA AGTCGAC 117

(2) INFORMATION FOR SEQ ID NO:17:
 50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Met Gly Arg Gly Asp
1 5

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GTCGACATGG GTCGTGGTGA TTAAGTCGAC 30

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Met Gly Arg Gly Asp Gly Lys Ser Gly Arg Gly Asp Gly Lys Ser Gly
1 5 10 15
Arg Gly Asp Gly Lys Ser
20

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 81 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GTCGACATGG GTCGTGGTGA TGGTAAAAGT GTCGTGGTG ATGGTAAAAG TGGTCGTGGT 60
GATGGTAAAA GTTAAGTCGA C 81

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

EP 0 921 193 A1

(A) LENGTH: 210 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

10

Met Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile Gly Asn Gly Tyr Arg
1 5 10 15

Gly Thr Met Ser Arg Thr Lys Ser Gly Val Ala Cys Gln Lys Trp Gly
20 25 30

15

Ala Thr Phe Pro His Val Pro Asn Tyr Ser Pro Ser Thr His Pro Asn
35 40 45

Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Glu Gln
50 55 60

20

Gly Pro Trp Cys Tyr Thr Thr Asp Pro Asp Lys Arg Tyr Asp Tyr Cys
65 70 75 80

Asn Ile Pro Glu Cys Glu Glu Glu Cys Met Tyr Cys Ser Gly Glu Lys
85 90 95

25

Tyr Glu Gly Lys Ile Ser Lys Thr Met Ser Gly Lys Asp Cys Gln Ala
100 105 110

Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ala Lys Phe
115 120 125

30

Pro Ser Lys Asn Leu Lys Met Asn Tyr Cys His Asn Pro Asp Gly Glu
130 135 140

Pro Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr Lys Arg Trp Glu Tyr
145 150 155 160

35

Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Pro Ser Pro Thr
165 170 175

Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser
180 185 190

40

Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro
195 200 205

His Arg
210

45

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

50

(A) LENGTH: 645 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

55

EP 0 921 193 A1

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:22:

5 GTCGACATGG TGTATCTGTC AGAATGTAAG ACCGGCATCG GCAACGGCTA CAGAGGAACC 60
 ATGTCCAGGA CAAAGAGTGG TGTTGCCTGT CAAAAGTGGG GTGCCACGTT CCCCCACGTA 120
 CCCAACTACT CTCCCAGTAC ACATCCCAAT GAGGGACTAG AAGAGAACTA CTGTAGGAAC 180
 CCAGACAATG ATGAACAAGG GCCTTGGTGC TAACTACAG ATCCGGACAA GAGATATGAC 240
 10 TACTGCAACA TTCCTGAATG TGAAGAGGAA TGCATGTACT GCAGTGGAGA AAAGTATGAG 300
 GGCAAAATCT CCAAGACCAT GTCTGGACTT GACTGCCAGG CCTGGGATTC TCAGAGCCCA 360
 CATGCTCATG GATACATCCC TGCCAAATTT CCAAGCAAGA ACCTGAAGAT GAATTATTGC 420
 15 CACAACCCTG ACGGGGAGCC AAGGCCCTGG TGCTTCACAA CAGACCCAC CAAACGCTGG 480
 GAATACTGTG ACATCCCCCG CTGCACAACA CCCCCGCCCC CACCCAGCCC AACCTACCAA 540
 TGTCTGAAAG GAAGAGGTGA AAATTACCGA GGGACCGTGT CTGTCACCGT GTCTGGGAAA 600
 20 ACCTGTCAGC GCTGGAGTGA GCAAACCCCT CATAGGTGAG TCGAC 644

(2) INFORMATION FOR SEQ ID NO:23:

(1) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 423 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Met Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile Gly Asn Gly Tyr Arg
 1 5 10 15
 35 Gly Thr Met Ser Arg Thr Lys Ser Gly Val Ala Cys Gln Lys Trp Gly
 20 25 30
 Ala Thr Phe Pro His Val Pro Asn Tyr Ser Pro Ser Thr His Pro Asn
 35 40 45
 40 Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Glu Gln
 50 55 60
 Gly Pro Trp Cys Tyr Thr Thr Asp Pro Asp Lys Arg Tyr Asp Tyr Cys
 65 70 75 80
 45 Asn Ile Pro Glu Cys Glu Glu Glu Cys Met Tyr Cys Ser Gly Glu Lys
 85 90 95
 Tyr Glu Gly Lys Ile Ser Lys Thr Met Ser Gly Lys Asp Cys Gln Ala
 100 105 110
 50 Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ala Lys Phe
 115 120 125
 Pro Ser Lys Asn Leu Lys Met Asn Tyr Cys His Asn Pro Asp Gly Glu

EP 0 921 193 A1

	130		135		140
5	Pro Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr Lys Arg Trp Glu Tyr				
	145		150		155
	Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Ser Pro Thr				
		165		170	175
10	Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser				
		180		185	190
	Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro				
		195		200	205
15	His Arg Gly Lys Ser Met Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile				
		210		215	220
	Gly Asn Gly Tyr Arg Gly Thr Met Ser Arg Thr Lys Ser Gly Val Ala				
		225		230	235
20	Cys Gln Lys Trp Gly Ala Thr Phe Pro His Val Pro Asn Tyr Ser Pro				
		245		250	255
	Ser Thr His Pro Asn Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro				
		260		265	270
25	Asp Asn Asp Glu Gln Gly Pro Trp Cys Tyr Thr Thr Asp Pro Asp Lys				
		275		280	285
	Arg Tyr Asp Tyr Cys Asn Ile Pro Glu Cys Glu Glu Glu Cys Met Tyr				
		290		295	300
30	Cys Ser Gly Glu Lys Tyr Glu Gly Lys Ile Ser Lys Thr Met Ser Gly				
		305		310	315
	Lys Asp Cys Gln Ala Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr				
		325		330	335
35	Ile Pro Ala Lys Phe Pro Ser Lys Asn Leu Lys Met Asn Tyr Cys His				
		340		345	350
	Asn Pro Asp Gly Glu Pro Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr				
		355		360	365
40	Lys Arg Trp Glu Tyr Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro				
		370		375	380
	Pro Pro Ser Pro Thr Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr				
		385		390	395
45	Arg Gly Thr Val Ser Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp				
		405		410	415
	Ser Glu Gln Thr Pro His Arg				
		420			

(2) INFORMATION FOR SEQ ID NO:24:

- (1) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1284 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

EP 0 921 193 A1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

	GTCGACATGG TGTATCTGTC AGAATGTAAG ACCGGCATCG GCAACGGCTA CAGAGGAACC	60
5	ATGTCCAGGA CAAAGAGTGG TGTTCCTGT CAAAAGTGGG GTGCCACGTT CCCCCACGTA	120
	CCCAACTACT CTCCCAGTAC ACATCCCAAT GAGGGACTAG AAGAGAACTA CTGTAGGAAC	180
	CCAGACAATG ATGAACAAGG GCCTTGGTGC TACACTACAG ATCCGGACAA GAGATATGAC	240
10	TACTGCAACA TTCCTGAATG TGAAGAGGAA TGCATGTACT GCAGTGGAGA AAAGTATGAG	300
	GGCAAAATCT CCAAGACCAT GTCTGGACTT GACTGCCAGG CCTGGGATTC TCAGAGCCCA	360
	CATGCTCATG GATACATCCC TGCCAAATTT CCAAGCAAGA ACCTGAAGAT GAATTATTGC	420
15	CACAACCCTG ACGGGGAGCC AAGGCCCTGG TGCTTCACAA CAGACCCAC CAAACGCTGG	480
	GAATACTGTG ACATCCCCCG CTGCACAACA CCCCCGCCCC CACCCAGCCC AACCTACCAA	540
	TGTCTGAAAG GAAGAGGTGA AAATTACCGA GGGACCGTGT CTGTCACCGT GTCTGGGAAA	600
20	ACCTGTCAGC GCTGGAGTGA GCAAACCCCT CATAGGGGTA AAAGTATGGT GTATCTGTCA	660
	GAATGTAAGA CCGGCATCGG CAACGGCTAC AGAGGAACCA TGTCCAGGAC AAAGAGTGGT	720
	GTTGCCTGTC AAAAGTGGGG TGCCACGTTC CCCCACGTAC CCAACTACTC TCCCAGTACA	780
25	CATCCCAATG AGGGACTAGA AGAGAACTAC TGTAGGAACC CAGACAATGA TGAACAAGGG	840
	CCTTGGTGCT ACACTACAGA TCCGGACAAG AGATATGACT ACTGCAACAT TCCTGAATGT	900
30	GAAGAGGAAT GCATGTACTG CAGTGGAGAA AAGTATGAGG GCAAAATCTC CAAGACCATG	960
	TCTGGACTTG ACTGCCAGGC CTGGGATTCT CAGAGCCAC ATGCTCATGG ATACATCCCT	1020
	GCCAAATTC CAAGCAAGAA CCTGAAGATG AATTATTGCC ACAACCCTGA CGGGGAGCCA	1080
35	AGGCCCTGGT GCTTCACAAC AGACCCACC AAACGCTGGG AATACTGTGA CATCCCCCGC	1140
	TGCACAACAC CCCCCCCCCC ACCCAGCCA ACCTACCAAT GTCTGAAAGG AAGAGGTGAA	1200
	AATTACCGAG GGACCGTGTG TGTCACCGTG TCTGGGAAAA CCTGTCAGCG CTGGAGTGAG	1260
40	CAAACCCCTC ATAGGTGAGT CGAC	1284

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 125 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

EP 0 921 193 A1

Met Leu Pro Ile Cys Pro Gly Gly Ala Ala Arg Cys Gln Val Thr Leu
1 5 10 15

Arg Glu Leu Phe Asp Arg Ala Val Val Leu Ser His Tyr Ile His Asn
5 20 25 30

Leu Ser Ser Glu Met Phe Ser Glu Phe Glu Lys Arg Tyr Thr His Gly
35 40 45

Arg Gly Phe Ile Thr Lys Ala Ile Asn Ser Cys His Thr Ser Ser Leu
10 50 55 60

Ala Thr Pro Glu Asp Lys Glu Gln Ala Gln Gln Met Asn Gln Lys Asp
65 70 75 80

Phe Leu Ser Leu Ile Val Ser Ile Leu Arg Ser Trp Asn Glu Pro Leu
15 85 90 95

Tyr His Leu Val Thr Glu Val Arg Gly Met Gln Glu Ala Pro Gln Ala
100 105 110

Ile Leu Ser Lys Ala Val Glu Ile Glu Glu Gln Thr Lys
20 115 120 125

(2) INFORMATION FOR SEQ ID NO:26:

(1) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 390 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:26:

GTGACATGT TGCCCATCTG TCCCGGCGGG GCTGCCCCGAT GCCAGGTGAC CCTTCGAGAC 60
CTGTTTGACC GCGCCGTCGT CCTGTCCCAC TACATCCATA ACCTCTCCTC AGAAATGTTC 120
AGCGAATTCG ATAAACGGTA TACCCATGGC CGGGGGTTCA TTACCAAGGC CATCAACAGC 180
TGCCCACTT CTTCCCTTGC CACCCCCGAA GACAAGGAGC AAGCCCAACA GATGAATCAA 240
AAAGACTTTC TGAGCCTGAT AGTCAGCATA TTGCGATCCT GGAATGAGCC TCTGTATCAT 300
CTGGTCACGG AAGTACGTGG TATGCAAGAA GCCCGGAGG CTATCCTATC CAAAGCTGTA 360
GAGATTGAGG AGCAAACCAA ATAAGTCGAC 390

45

(2) INFORMATION FOR SEQ ID NO:27:

(1) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 253 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

50

55

EP 0 921 193 A1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

5	Met	Leu	Pro	Ile	Cys	Pro	Gly	Gly	Ala	Ala	Arg	Cys	Gln	Val	Thr	Leu	1	5	10	15
	Arg	Glu	Leu	Phe	Asp	Arg	Ala	Val	Val	Leu	Ser	His	Tyr	Ile	His	Asn	20	25	30	
10	Leu	Ser	Ser	Glu	Met	Phe	Ser	Glu	Phe	Glu	Lys	Arg	Tyr	Thr	His	Gly	35	40	45	
	Arg	Gly	Phe	Ile	Thr	Lys	Ala	Ile	Asn	Ser	Cys	His	Thr	Ser	Ser	Leu	50	55	60	
15	Ala	Thr	Pro	Glu	Asp	Lys	Glu	Gln	Ala	Gln	Gln	Met	Asn	Gln	Lys	Asp	65	70	75	80
	Phe	Leu	Ser	Leu	Ile	Val	Ser	Ile	Leu	Arg	Ser	Trp	Asn	Glu	Pro	Leu	85	90	95	
20	Tyr	His	Leu	Val	Thr	Glu	Val	Arg	Gly	Met	Gln	Glu	Ala	Pro	Gln	Ala	100	105	110	
	Ile	Leu	Ser	Lys	Ala	Val	Glu	Ile	Glu	Glu	Gln	Thr	Lys	Gly	Lys	Ser	115	120	125	
25	Met	Leu	Pro	Ile	Cys	Pro	Gly	Gly	Ala	Ala	Arg	Cys	Gln	Val	Thr	Leu	130	135	140	
	Arg	Glu	Leu	Phe	Asp	Arg	Ala	Val	Val	Leu	Ser	His	Tyr	Ile	His	Asn	145	150	155	160
30	Leu	Ser	Ser	Glu	Met	Phe	Ser	Glu	Phe	Glu	Lys	Arg	Tyr	Thr	His	Gly	165	170	175	
	Arg	Gly	Phe	Ile	Thr	Lys	Ala	Ile	Asn	Ser	Cys	His	Thr	Ser	Ser	Leu	180	185	190	
35	Ala	Thr	Pro	Glu	Asp	Lys	Glu	Gln	Ala	Gln	Gln	Met	Asn	Gln	Lys	Asp	195	200	205	
	Phe	Leu	Ser	Leu	Ile	Val	Ser	Ile	Leu	Arg	Ser	Trp	Asn	Glu	Pro	Leu	210	215	220	
40	Tyr	His	Leu	Val	Thr	Glu	Val	Arg	Gly	Met	Gln	Glu	Ala	Pro	Gln	Ala	225	230	235	240
45	Ile	Leu	Ser	Lys	Ala	Val	Glu	Ile	Glu	Glu	Gln	Thr	Lys				245	250		

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 771 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

EP 0 921 193 A1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

5 GTCGACATGT TGCCCATCTG TCCCGGCGGG GCTGCCCGAT GCCAGGTGAC CCTTCGAGAC 60
CTGTTTGACC GCGCCGTCGT CCTGTCCAC TACATCCATA ACCTCTCCTC AGAAATGTTT 120
AGCGAATTCG ATAAACGGTA TACCCATGGC CGGGGGTTCA TTACCAAGGC CATCAACAGC 180
TGCCACACTT CTTCCCTTGC CACCCCGAA GACAAGGAGC AAGCCCAACA GATGAATCAA 240
10 AAAGACTTTC TGAGCCTGAT AGTCAGCATA TTGCGATCCT GGAATGAGCC TCTGTATCAT 300
CTGGTCACGG AAGTACGTGG TATGCAAGAA GCGCCGAGG CTATCCTATC CAAAGCTGTA 360
GAGATTGAGG AGCAAACCGG TAAAGTATG TTGCCCATCT GTCCCGCGGG GGCTGCCCGA 420
15 TGCCAGGTGA CCCTTCGAGA CCTGTTTGAC CGCGCCGTCG TCCTGTCCCA CTACATCCAT 480
AACCTCTCCT CAGAAATGTT CAGCGAATTC GATAACGGT ATACCCATGG CCGGGGGTTC 540
ATTACCAAGG CCATCAACAG CTGCCACACT TCTTCCCTTG CCACCCCGA AGACAAGGAG 600
20 CAAGCCCAAC AGATGAATCA AAAAGACTTT CTGAGCCTGA TAGTCAGCAT ATTGCGATCC 660
TGGAATGAGC CTCTGTATCA TCTGGTCACG GAAGTACGTG GTATGCAAGA AGCCCGGAG 720
GCTATCCTAT CCAAAGCTGT AGAGATTGAG GAGCAAACCA AATAAGTCGA C 771

25 (2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 161 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
30 (D) TOPOLOGY: linear

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

ATGCTGAGGC GGCCTCCCCT ATGCTATCAC AACGGAGTTC AGTACAGAAA TAACGGTAAA 60
AGATCCCCGT GGTCATCTTG TTCTGTGACA TGTGGTGATG GTGTGATGGT AAAAGAAGTG 120
40 GTACCCTGTA GACAAGACAG TGGACACCTC CTCCCCATTA A 161

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 63 amino acids
(B) TYPE: amino acid
45 (D) TOPOLOGY: linear

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Met Leu Arg Arg Pro Pro Leu Cys Tyr His Asn Gly Val Gln Tyr Arg
1 5 10 15

55

EP 0 921 193 A1

Asn Asn Glu Glu Trp Thr Val Asp Ser Gly Lys Ser Ser Pro Trp Ser
 20 25 30
 Ser Cys Ser Val Thr Cys Gly Asp Gly Val Ile Thr Arg Ile Gly Lys
 35 40 45
 Ser Ser Pro Trp Asp Ile Cys Ser Val Thr Cys Gly Gly Gly Val
 50 55 60

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 185 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

ATGCTGAGGC GGCCTCCCCCT ATGCTATCAC AACGGAGTTC AGTACAGAAA TAACGGTAAA 60
 AGATCCCCGT GGTTCATCTTG TTCTGTGACA TGTGGTGATG GTGTGATGGT AAAAGAAGTG 120
 GTACCCTGTA GACAAGACAG TGGACACCTC CTCCCCATTA TATTGGTTCT CGTGGTAAAA 180
 GATAA 185

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

TAGGTCTAGA ATGACTGAAG AGAACAAAGA G 31

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

ATGGTCTAGA TTAGAGACGA CTACGTTTCT G 31

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 805 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Met Glu Ser Lys Ala Leu Leu Ala Val Ala Leu Trp Phe Cys Val Glu
 1 5 10 15

Thr Arg Ala Ala Ser Val Gly Leu Pro Gly Asp Phe Leu His Pro Pro
 20 25 30

Lys Leu Ser Thr Gln Lys Asp Ile Leu Thr Ile Leu Ala Asn Thr Thr
 35 40 45

Leu Gln Ile Thr Cys Arg Gly Gln Arg Asp Leu Asp Trp Leu Trp Pro
 50 55 60

Asn Ala Gln Arg Asp Ser Glu Glu Arg Val Leu Val Thr Glu Cys Gly
 65 70 75 80

Gly Gly Asp Ser Ile Phe Cys Lys Thr Leu Thr Ile Pro Arg Val Val
 85 90 95

Gly Asn Asp Thr Gly Ala Tyr Lys Cys Ser Tyr Arg Asp Val Asp Ile
 100 105 110

Ala Ser Thr Val Tyr Val Tyr Val Arg Asp Tyr Arg Ser Pro Phe Ile
 115 120 125

Ala Ser Val Ser Asp Gln His Gly Ile Val Tyr Ile Thr Glu Asn Lys
 130 135 140

Asn Lys Thr Val Val Ile Pro Cys Arg Gly Ser Ile Ser Asn Leu Asn
 145 150 155 160

Val Ser Leu Cys Ala Arg Tyr Pro Glu Lys Arg Phe Val Pro Asp Gly
 165 170 175

Asn Arg Ile Ser Trp Asp Ser Glu Ile Gly Phe Thr Leu Pro Ser Tyr
 180 185 190

Met Ile Ser Tyr Ala Gly Met Val Phe Cys Glu Ala Lys Ile Asn Asp
 195 200 205

Glu Thr Tyr Gln Ser Ile Met Tyr Ile Val Val Val Val Gly Tyr Arg
 210 215 220

Ile Tyr Asp Val Ile Leu Ser Pro Pro His Glu Ile Glu Leu Ser Ala
 225 230 235 240

Gly Glu Lys Leu Val Leu Asn Cys Thr Ala Arg Thr Glu Leu Asn Val
 245 250 255

Gly Leu Asp Phe Thr Trp His Ser Pro Pro Ser Lys Ser His His Lys

EP 0 921 193 A1

	260	265	270
	Lys Ile Val Asn Arg Asp Val	Lys Pro Phe Pro Gly Thr Val Ala Lys	
5	275	280	285
	Met Phe Lys Ser Thr Leu Thr	Ile Glu Ser Val Thr Lys Ser Asp Gln	
	290	295	300
10	Gly Glu Tyr Thr Cys Val Ala Ser Ser Gly Arg Met Ile Lys Arg Asn		
	305	310	315
	Arg Thr Phe Val Arg Val His Thr Lys Pro Phe Ile Ala Phe Gly Ser		
	325	330	335
15	Gly Met Lys Ser Leu Val Glu Ala Thr Val Gly Ser Gln Val Arg Ile		
	340	345	350
	Pro Val Lys Tyr Leu Ser Tyr Pro Ala Pro Asp Ile Lys Trp Tyr Arg		
	355	360	365
20	Asn Gly Arg Pro Ile Glu Ser Asn Tyr Thr Met Ile Val Gly Asp Glu		
	370	375	380
	Leu Thr Ile Met Glu Val Thr Glu Arg Asp Ala Gly Asn Tyr Thr Val		
	385	390	395
25	Ile Leu Thr Asn Pro Ile Ser Met Glu Lys Gln Ser His Met Val Ser		
	405	410	415
	Leu Val Val Asn Val Pro Pro Gln Ile Gly Glu Lys Ala Leu Ile Ser		
	420	425	430
30	Pro Met Asp Ser Tyr Gly Tyr Gly Thr Met Gln Thr Leu Thr Cys Thr		
	435	440	445
	Val Tyr Ala Asn Pro Pro Leu His His Ile Gln Trp Tyr Trp Gln Leu		
	450	455	460
35	Glu Glu Ala Cys Ser Tyr Arg Pro Gly Gln Thr Ser Pro Tyr Ala Cys		
	465	470	475
	Lys Glu Trp Arg His Val Glu Asp Phe Gln Gly Gly Asn Lys Ile Glu		
	485	490	495
40	Val Thr Lys Asn Gln Tyr Ala Leu Ile Glu Gly Lys Asn Lys Thr Val		
	500	505	510
	Ser Thr Leu Val Ile Gln Ala Ala Asn Val Ser Ala Leu Tyr Lys Cys		
	515	520	525
45	Glu Ala Ile Asn Lys Ala Gly Arg Gly Glu Arg Val Ile Ser Phe His		
	530	535	540
	Val Ile Arg Gly Pro Glu Ile Thr Val Gln Pro Ala Ala Gln Pro Thr		
	545	550	555
50	Glu Gln Glu Ser Val Ser Leu Leu Cys Thr Ala Asp Arg Asn Thr Phe		
	565	570	575
	Glu Asn Leu Thr Trp Tyr Lys Leu Gly Ser Gln Ala Thr Ser Val His		
	580	585	590

EP 0 921 193 A1

	Met	Gly	Glu	Ser	Leu	Thr	Pro	Val	Cys	Lys	Asn	Leu	Asp	Ala	Leu	Trp
			595					600					605			
5	Lys	Leu	Asn	Gly	Thr	Met	Phe	Ser	Asn	Ser	Thr	Asn	Asp	Ile	Leu	Ile
		610					615					620				
	Val	Ala	Phe	Gln	Asn	Ala	Ser	Leu	Gln	Asp	Gln	Gly	Asp	Tyr	Val	Cys
	625					630					635					640
10	Ser	Ala	Gln	Asp	Lys	Lys	Thr	Lys	Lys	Arg	His	Cys	Leu	Val	Lys	Gln
					645					650					655	
	Leu	Ile	Ile	Leu	Glu	Arg	Met	Ala	Pro	Met	Ile	Thr	Gly	Asn	Leu	Glu
				660					665					670		
15	Asn	Gln	Thr	Thr	Thr	Ile	Gly	Glu	Thr	Ile	Glu	Val	Thr	Cys	Pro	Ala
				675				680						685		
	Ser	Gly	Asn	Pro	Thr	Pro	His	Ile	Thr	Trp	Phe	Lys	Asp	Asn	Glu	Thr
		690					695					700				
20	Leu	Val	Glu	Asp	Ser	Gly	Ile	Val	Leu	Arg	Asp	Gly	Asn	Arg	Asn	Leu
	705					710					715					720
	Thr	Ile	Arg	Arg	Val	Arg	Lys	Glu	Asp	Gly	Gly	Leu	Tyr	Thr	Cys	Gln
					725					730					735	
25	Ala	Cys	Asn	Val	Leu	Gly	Cys	Ala	Arg	Ala	Glu	Thr	Leu	Phe	Ile	Ile
				740					745					750		
	Glu	Gly	Ala	Gln	Glu	Lys	Thr	Asn	Leu	Glu	Val	Ile	Ile	Leu	Val	Gly
			755					760					765			
30	Thr	Ala	Val	Ile	Ala	Met	Phe	Phe	Trp	Leu	Leu	Leu	Val	Ile	Leu	Val
		770					775					780				
	Arg	Thr	Val	Lys	Arg	Ala	Asn	Glu	Gly	Glu	Leu	Lys	Thr	Gly	Tyr	Leu
	785					790					795					800
35	Ser	Ile	Val	Met	Asp											
				805												

(2) INFORMATION FOR SEQ ID NO:35:

	(i) SEQUENCE CHARACTERISTICS:
40	(A) LENGTH: 2431 base pairs
	(B) TYPE: nucleic acid
	(C) STRANDEDNESS: single
	(D) TOPOLOGY: linear

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

	AGACGTCATG GAGAGCAAGG CGCTGCTAGC TGTCGCTCTG TGGTTCTGCG TGGAGACCCG	60
50	AGCCGCCTCT GTGGGTTTGC CTGGCGATTT TCTCCATCCC CCCAAGCTCA GCACACAGAA	120
	AGACATACTG ACAATTTTGG CAAATACAAC CCTTCAGATT ACTTGCAGGG GACAGCGGGA	180

55

EP 0 921 193 A1

	CCTGGACTGG	CTTTGGCCCA	ATGCTCAGCG	TGATTCTGAG	GAAAGGGTAT	TGGTGA	240
	ATGCGGCGGT	GGTGACAGTA	TCTTCTGCAA	AACACTCACC	ATTCCCAGGG	TGGTTGAAA	300
5	TGATACTGGA	GCCTACAAGT	GCTCGTACCG	GGACGTCGAC	ATAGCCTCCA	CTGTTTATGT	360
	CTATGTTTGA	GATTACAGAT	CACCATTTCAT	CGCCTCTGTC	AGTGACCAGC	ATGGCATCGT	420
	GTACATCACC	GAGAACAAGA	ACAAAACACTGT	GGTGATCCCC	TGCCGAGGGT	CGATTTCAAA	480
10	CCTCAATGTG	TCTCTTTGCG	CTAGGTATCC	AGAAAAGAGA	TTTGTTCCGG	ATGGAAACAG	540
	AATTTCTTGG	GACAGCGAGA	TAGGCTTTAC	TCTCCCCAGT	TACATGATCA	GCTATGCCGG	600
	CATGGTCTTC	TGTGAGGCAA	AGATCAATGA	TGAAACCTAT	CAGTCTATCA	TGTACATAGT	660
15	TGTGGTTGTA	GGATATAGGA	TTTATGATGT	GATTCTGAGC	CCCCCGCATG	AAATTGAGCT	720
	ATCTGCCGGA	GAAAACTTG	TCTTAAATTG	TACAGCGAGA	ACAGAGCTCA	ATGTGGGGCT	780
	TGATTTCAAC	TGGCACTCTC	CACCTTCAAA	GTCTCATCAT	AAGAAGATTG	TAAACCGGGA	840
20	TGTGAAACCC	TTTCTTGGGA	CTGTGGCGAA	GATGTTTTTG	AGCACCTTGA	CAATAGAAAG	900
	TGTGACCAAG	AGTGACCAAG	GGGAATACAC	CTGTGTAGCG	TCCAGTGGAC	GGATGATCAA	960
	GAGAAATAGA	ACATTTGTCC	GAGTTCACAC	AAAGCCTTTT	ATTGCTTTTCG	GTAGTGGGAT	1020
25	GAAATCTTTG	GTGGAAGCCA	CAGTGGGCAG	TCAAGTCCGA	ATCCCTGTGA	AGTATCTCAG	1080
	TTACCCAGCT	CCTGATATCA	AATGGTACAG	AAATGGAAGG	CCCATTGAGT	CCAACTACAC	1140
	AATGATTGTT	GGCGATGAAC	TCACCATCAT	GGAAGTGA	GAAAGAGATG	CAGGAAACTA	1200
30	CACGGTCATC	CTCACCAACC	CCATTTCAAT	GGAGAAACAG	AGCCACATGG	TCTCTCTGGT	1260
	TGTGAATGTC	CCACCCCAAG	TCGGTGAGAA	AGCCTTGATC	TCGCCTATGG	ATTCCTACCA	1320
	GTATGGGACC	ATGCAGACAT	TGACATGCAC	AGTCTACGCC	AACCCCTCCC	TGCACCACAT	1380
35	CCAGTGGTAC	TGGCAGCTAG	AAGAAGCCTG	CTCCTACAGA	CCCGGCCAAA	CAAGCCCGTA	1440
	TGCTTGTA	GAATGGAGAC	ACGTGGAGGA	TTTCCAGGGG	GGAAACAAGA	TCGAAGTCAC	1500
	CAAAACCAA	TATGCCCTGA	TTGAAGGAAA	AAACAAACT	GTAAGTACGC	TGGTCATCCA	1560
40	AGCTGCCAAC	GTGTCAGCGT	TGTACAAATG	TGAAGCCATC	AACAAAGCGG	GACGAGGAGA	1620
	GAGGGTCATC	TCCTTCCATG	TGATCAGGGG	TCCTGAAATT	ACTGTGCAAC	CTGCTGCCCA	1680
	GCCAACTGAG	CAGGAGAGTG	TGTCCCTGTT	GTGCACTGCA	GACAGAAATA	CGTTTGAGAA	1740
45	CCTCACGTGG	TACAAGCTTG	GCTCACAGGC	AACATCGGTC	CACATGGGCG	AATCACTCAC	1800
	ACCAGTTTGC	AAGAACTTGG	ATGCTCTTTG	GAAACTGAAT	GGCACCATGT	TTTCTAACAG	1860
	CACAAATGAC	ATCTTGATTG	TGGCATTTC	GAATGCCTCT	CTGCAGGACC	AAGGCGACTA	1920
50	TGTTTGCTCT	GCTCAAGATA	AGAAGACCAA	GAAAAGACAT	TGCCTGGTCA	AACAGCTCAT	1980
	CATCCTAGAG	CGCATGGCAC	CCATGATCAC	CGGAAATCTG	GAGAATCAGA	CAACAACCAT	2040

55

EP 0 921 193 A1

TGGCGAGACC ATTGAAGTGA CTTGCCCAGC ATCTGGAAAT CCTACCCAC ACATTACATG 2100
 GTTCAAAGAC AACGAGACCC TGGTAGAAGA TTCAGGCATT GTACTGAGAG ATGGGAACCG 2160
 5 GAACCTGACT ATCCGCAGGG TGAGGAAGGA GGATGGAGGC CTCTACACCT GCCAGGCCTG 2220
 CAATGTCCTT GGCTGTGCAA GAGCGGAGAC GCTCTTCATA ATAGAAGGTG CCCAGGAAAA 2280
 GACCAACTTG GAAGTCATTA TCCTCGTCGG CACTGCAGTG ATTGCCATGT TCTTCTGGCT 2340
 10 CCTTCTTGTC ATTCTCGTAC GGACCGTTAA GCGGGCCAAT GAAGGGGAAC TGAAGACAGG 2400
 CTACTTGTCT ATTGTCATGG ATTAAGACGT C 2431

(2) INFORMATION FOR SEQ ID NO:36:

15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 185 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

25 Met His Thr His Gln Asp Phe Gln Pro Val Leu His Leu Val Ala Leu
 1 5 10 15
 Asn Thr Pro Leu Ser Gly Gly Met Arg Gly Ile Arg Gly Ala Asp Phe
 20 25 30
 30 Gln Cys Phe Asn Asn Ala Arg Val Gly Leu Ser Gly Thr Phe Arg Ala
 35 40 45
 Phe Leu Ser Ser Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg Arg Ala
 50 55 60
 35 Asp Arg Gly Ser Val Pro Ile Val Gln Asn Leu Arg Asp Glu Val Leu
 65 70 75 80
 Ser Pro Ser Trp Asp Ser Leu Phe Ser Gly Ser Gln Gly Gln Leu Gln
 85 90 95
 40 Pro Gly Ala Arg Ile Phe Ser Phe Asp Gly Arg Asp Val Leu Arg His
 100 105 110
 Pro Ala Trp Pro Gln Arg Ser Val Trp His Gly Ser Asp Pro Ser Gly
 115 120 125
 45 Arg Arg Leu Met Glu Ser Tyr Cys Glu Thr Trp Arg Thr Glu Thr Thr
 130 135 140
 Gly Ala Thr Gly Gln Ala Ser Ser Leu Leu Ser Gly Arg Leu Leu Glu
 145 150 155 160
 50 Gln Arg Ala Ala Ser Cys His Asp Ser Tyr Ile Val Leu Cys Ile Glu
 165 170 175
 Asn Ser Phe Met Thr Ser Phe Ser Arg
 180 185

55

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 565 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

AGACGTCATG CATACTCATC AGGACTTTCA GCCAGTGCTC CACCTGGTGG CACTGAACAC 60
 CCCCCTGTCT GGAGGCATGC GTGGTATCCG TGGAGCAGAT TTCCAGTGCT TCCAGCAAGC 120
 CCGAGCCGTG GGGCTGTCGG GCACCTTCCG GGCTTTCCTG TCCTCTAGGC TGCAGGATCT 180
 CTATAGCATC GTGCGCCGTG CTGACCGGGG GTCTGTGCCC ATCGTCAACC TGAAGGACGA 240
 GGTGCTATCT CCCAGCTGGG ACTCCCTGTT TTCTGGCTCC CAGGGTCAAC TGCAACCCGG 300
 GGCCCGCATC TTTTCTTTTG ACGGCAGAGA TGTCTGAGA CACCCAGCCT GGCCGCAGAA 360
 GAGCGTATGG CACGGCTCGG ACCCCAGTGG GCGGAGGCTG ATGGAGAGTT ACTGTGAGAC 420
 ATGGCGAACT GAAACTACTG GGGCTACAGG TCAGGCCTCC TCCCTGCTGT CAGGCAGGCT 480
 CCTGGAACAG AAAGCTGCGA GCTGCCACAA CAGCTACATC GTCCTGTGCA TTGAGAATAG 540
 CTTTCATGACC TCTTCTCCA AATAG 565

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 43 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

CTATCGTCGA CATGTATATT GGTTCCTCGTT AAGTCGACCT ATC 43

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 43 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

GATAGGTCGA CTTAACGAGA ACCAATATAC ATGTCGACGA TAG

43

(2) INFORMATION FOR SEQ ID NO:40:

5

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

15

AGTATCTAGA ATGAGTGTAT CTGTCACAAT G

31

(2) INFORMATION FOR SEQ ID NO:41:

20

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

30

GAATTCTAGA TCACCTATGA GGGGTTTGCT C

31

(2) INFORMATION FOR SEQ ID NO:42:

35

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 93 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

CTATCGTCGA CATGTATATT GGTTCGTA AAAGATATAT TGGTTCTCGT GGTAAAAGAG

60

45

ATGGTTCTCG TGGTAAAGA TAAGTGACCT ATC

93

(2) INFORMATION FOR SEQ ID NO:43:

50

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

55

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:43:

GATAGGTCGA CTTAT


15

5


Claims

- 10 1. A carrier:DNA complex comprising DNA encoding at least one anti-angiogenic protein or peptide, the complex being deliverable to the site of a tumor in vivo and additionally comprising regulatory elements for expressing the anti-angiogenic DNA in a tumor or tumor vasculature.
- 15 2. The complex of claim 1, wherein the carrier is selected from the group consisting of liposomes, cationic polymers, micelles, microspheres, viruses, viral components, or combinations of such carriers.
3. The complex of claim 1, wherein the complex additionally contains DNA encoding a tumor suppressor protein.
- 20 4. The complex of Claim 2, wherein the complex additionally contains DNA encoding a tumor suppressor protein.
5. The complex of claim 3 or 4, wherein the tumor suppressor protein is p53.
6. The complex of claim 1, additionally comprising a marker directing the complexes in vivo to a tumor or to tumor or peritumoral area.
- 25 7. The complex of claim 1, wherein the DNA is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25; SEQ ID NO:27, SEQ ID NO:31; SEQ ID NO:35; and SEQ ID NO: 37.
- 30 8. The complex of claim 1, wherein the anti-angiogenic DNA is provided in a vector containing at least one promotor.
- 35 9. Use of DNA encoding at least one anti-angiogenic protein or peptide in a carrier preferably as defined in any of claims 1 to 8 which is delivered to a tumor site in vivo and is expressed at the tumor site in vivo in the subject for the production of a medicament for inhibiting tumor growth in a subject bearing a tumor, which comprises administering the same to the subject.
- 40 10. The use of claim 9, wherein the carrier is selected from the group consisting of liposomes, cationic polymers, micelles, microspheres, viruses, viral components, or combinations of such carriers.
11. The use of claim 9, which further comprises providing a DNA encoding a tumor suppressor protein on the carrier.
- 45 12. The use of any of claims 9 to 11 wherein the administration is by injection.
13. The use of claim 9 to 11 wherein the administration is by intravenous injection.
- 50 14. Use of an anti-angiogenic DNA in a form in which the DNA is expressed in the tumor or a peritumoral area inhibiting tumor growth in a subject bearing a tumor, which comprises injecting the same into the subject.
- 55 15. The use of claim 14, wherein DNA encoding a tumor suppressor protein is additionally injected in a form which is expressed in the tumor or associated tumor vasculature.
16. The use of claim 14, wherein the injection is intravenous.
17. The use of claim 14, wherein the injection is intratumoral.

Figure 1
Intratumoral Injections of Liposome:DNA Complexes



*, Angio vs. BAP, $p < 0.05$



*- BAP vs BAP-p53 or BAP-TSPf, $p < 0.05$

** - BAP-p53 or BAP-TSPf vs BAP-p53/BAP-TSPf, $p < 0.01$



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 10 0135

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
E	EP 0 819 758 A (MIXSON, ARCHIBALD JAMES) 21 January 1998 * page 5, line 5 - line 54 * * page 13, line 46 - page 15, line 43 * * examples 2,3 * ---	1-17	C12N15/12 C12N15/24 C12N15/88 A61K48/00
X	WO 96 21416 A (CHIRON VIAGENE, INC.) 18 July 1996	1,2,6, 8-10, 12-14	
Y	* page 4, line 17 - line 22 * * page 5, line 7 - line 16 * * page 9, line 7 - page 10, line 23 * * page 15, line 8 - page 17, line 5 * * page 21, line 25 - page 23, line 9 * * page 30, line 12 - page 31, line 6 * ---	3-5,11, 15-17	
X	WO 95 29242 A (THE CHILDREN'S MEDICAL CENTER CORPORATION) 2 November 1995 * page 7, line 24 - page 8, line 34 * * page 16, line 15 - page 17, line 5 * * page 20, line 29 - page 24, line 2 * * page 26, line 13 - page 28, line 4 * * examples 23,24 * ---	1,2,6, 8-10,12, 14	TECHNICAL FIELDS SEARCHED (Int.Cl.6) A61K
Y	XU, M. ET AL.: "Parenteral gene therapy with p53 inhibits human breast tumors in vivo through a bystander mechanism without evidence of toxicity" HUMAN GENE THERAPY, vol. 8, no. 2, 20 January 1997, pages 177-185, XP002070390 * page 177 * *Abstract* * page 182 - page 183 * *Discussion* --- -/--	1-17	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 3 July 1998	Examiner Donath, C
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03/82 (P04C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 10 0135

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Y	<p>LESOON-WOOD, L.A. ET AL.: "Systemic gene therapy with p53 reduces growth and metastases of a malignant human breast cancer in nude mice"</p> <p>HUMAN GENE THERAPY, vol. 6, no. 4, April 1995, pages 395-405, XP002070391</p> <p>* page 395 *</p> <p>*Abstract*</p> <p>* page 401 - page 403 *</p> <p>*Discussion*</p>	1-17	
A,D	<p>WEINSTAT-SASLOW, D.L. ET AL.: "Transfection of thrombospondin 1 complementary DNA into a human breast carcinoma cell line reduces primary tumor growth, metastatic potential, and angiogenesis"</p> <p>CANCER RESEARCH, vol. 54, 15 December 1994, pages 6504-6511, XP002048545</p> <p>* page 6504 *</p> <p>*Abstract*</p> <p>* page 6508 - page 6510 *</p> <p>*Discussion*</p>	1-17	
The present search report has been drawn up for all claims			<p>TECHNICAL FIELDS SEARCHED (Int.Cl.6)</p>
Place of search		Date of completion of the search	Examiner
MUNICH		3 July 1998	Donath, C
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone</p> <p>Y : particularly relevant if combined with another document of the same category</p> <p>A : technological background</p> <p>O : non-written disclosure</p> <p>P : intermediate document</p> <p>T : theory or principle underlying the invention</p> <p>E : earlier patent document, but published on, or after the filing date</p> <p>D : document cited in the application</p> <p>L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 10 0135

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

03-07-1998

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0819758 A	21-01-1998	NONE	
WO 9621416 A	18-07-1996	AU 4608296 A EP 0802801 A	31-07-1996 29-10-1997
WO 9529242 A	02-11-1995	US 5639725 A AU 2461795 A BR 9507479 A CA 2188813 A CN 1149319 A CZ 9603122 A EP 0758390 A HU 76095 A JP 9512173 T US 5733876 A ZA 9503419 A	17-06-1997 16-11-1995 16-09-1997 02-11-1995 07-05-1997 14-05-1997 19-02-1997 30-06-1997 09-12-1997 31-03-1998 11-01-1996